


2015 specification
first exams in 2017



Titrations 'Tricky Topics' **Worksheets**

for A Level OCR Chemistry A

**Exam
skills prep
pack**

zigzageducation.co.uk

**POD
9444**

Publish your own work... Write to a brief...
Register at publishmenow.co.uk

Contents

Thank You for Choosing ZigZag Education.....	ii
Teacher Feedback Opportunity.....	iii
Terms and Conditions of Use	iv
Teacher's Introduction	1
Student Introduction	2
Worksheets.....	3
Titration basics: results and uncertainties	3
Concentrations and dilution	6
Structured acid–base titration calculations	8
Unstructured acid–base titration calculations	11
‘Back’ titrations	13
Titration curves / indicators	16
Strong acids and bases	20
Weak acids	22
Buffers.....	24
Structured redox titrations.....	26
Unstructured redox titrations.....	30
The most difficult redox titrations	33
Exam-style questions	37
pH Titrations	37
Redox Titrations	40
Appendix: GCSE knowledge refresher.....	42
Acids and alkalis	42
Concentrations and mole calculations in reactions	44
Reduction and oxidation	46
Quick quiz.....	47
Answers	48

Teacher's Introduction

This activity pack is designed to help your students develop skills both during class and in an extracurricular setting. The content covers titration where it occurs in a range of topics across the A Level course.

The resource contains:

- a student introduction
- 12 worksheets, each of which covers one of the skills outlined below
- exam-style questions
- an appendix – a refresher for GCSE topics relating to titrations

Although each worksheet covers a specific section of the A Level resource, the exercises are primarily skills-focused. These skills are all relevant to the A Level OCR A examinations and are covered as follows:

- | | |
|--|---|
| 1. Titration basics: results and uncertainties | 7. Strong acids and bases |
| 2. Concentrations and dilutions | 8. Weak acids |
| 3. Structured acid–base titration questions | 9. Buffers |
| 4. Unstructured acid–base titration questions | 10. Structured redox titrations |
| 5. Back titrations | 11. Unstructured redox titrations |
| 6. Titration curves and choosing indicators | 12. The most difficult redox titrations |

Each worksheet contains a short section of background information, followed by **worked examples** and then **in-depth questions** to test students' knowledge. The questions are split into **three levels of increasing difficulty**, illustrated using the headings *Prepare your solutions*, *High concentration* and *Drop the base*. You'll find an answers section at the end of the pack.

The pack also includes **full exam-style questions** which closely replicate the style of questions found in an examination paper.

I hope this resource will be useful to your teaching and will help your students not only to tackle an area of chemistry which many find challenging, but also to gain a deeper, more holistic understanding of the topic.

Free Updates!

Register your email address to receive any future free updates* made to this resource or other Chemistry resources your school has purchased, and details of any promotions for your subject.

* resulting from minor specification changes, suggestions from teachers and peer reviews, or occasional errors reported by customers

Go to [zzed.uk/freeupdates](https://www.zzed.uk/freeupdates)

Student Introduction

Acids



Sour-tasting substances were called 'oxein' by the ancient Greeks. The Greeks didn't know what was responsible for sour tastes were acids. With the Romans, this word became 'acetum' which is vinegar, which tastes sour due to the presence of ethanoic acid. It was also known that acids could change the colour of litmus, a type of moss, and that they could react with metals.

'Acetum' in Latin became 'acid' in English. This word now describes not only the acid in vinegar and citric acid in lemons, but also the acids you'd be better off not touching, like sulfuric acid, for example.

Arrhenius was first to propose that acids, when dissolved in water, produced H^+ ions. He also noted that alkalis produce solutions of OH^- ions, and that acids and alkalis react in water. We still use these definitions today.

Titration

Titration was first used in eighteenth-century France. Joseph Louis Gay-Lussac designed the side arm, and inexplicably making it so the tap stuck out on the wrong side. Below was another scientist, Biot, in a hot-air balloon in 1814, collecting gases at different altitudes. Biot improved the burette, adding a clamp and tip at the bottom, and then drawing some of the day, the sidearm tap of the burette remains curiously uncomfortable to reach even for chemists.



Purpose of this pack

Titration is probably among the more difficult practicals you'll do at school. You have to mix the solutions; swirling the flask and operating the tap at the same time always seem to require two hands, and even if you do manage to reach the end without your burette coming out, you've got to have a pretty good handle on the maths to make it all worth it.

This pack will equip you to deal with the really mathsy bits, starting at the basics, through exam questions, and everything you need to know in between:

1. Results tables and uncertainties
2. Concentrations and dilution
3. Structured acid-base titration questions
4. Unstructured acid-base titration questions
5. Back titration
6. Titration curves and choosing indicators
7. Strong acids and bases
8. Weak acids
9. Buffers
10. Structured redox titration questions
11. Unstructured redox titration questions
12. The most difficult redox titration questions

INSPECTION COPY

COPYRIGHT
PROTECTED




Titration basics: results and uncertainty

Key skills

- Reading a burette
- Completing titration results tables
- Calculating mean titres
- Calculating uncertainties

concordant – titrations, results
titre – the result
uncertainty –

Reading the burette



The reading from the bottom of the meniscus is given to 2 d.p.
 This burette reading is 19.60 cm³ and

Results tables

Usually, the first titre will be done very quickly to find a rough result. This result is not included in the mean.

After that, the titration is repeated until two concordant results are obtained. This is within 0.1 cm³ of each other.

An average of concordant results is then taken.

$$\text{mean titre} = \frac{\text{sum of all concordant titres}}{\text{number of concordant titres}}$$

Example table

	Rough	1	2	3
Start volume (cm³)	0.00	0.05	0.00	0.00
End volume (cm³)	25.65	23.30	23.50	23.35
Volume of acid (cm³)	25.65	23.25	23.50	23.35

The experiment is repeated to increase accuracy.

End volume – Start volume

Titre 2 is **not concordant** with titres 1 and 3, so it is not included in the mean.
 Titres 1 and 3 are **concordant** – they are within 0.10 cm³ of each other.

COPYRIGHT
PROTECTED



Uncertainties

Titration results come with a certain amount of **uncertainty**. The main factor in this uncertainty is the uncertainty due to the burette.

The uncertainty of a burette is $\pm 0.05 \text{ cm}^3$ per reading. There are two readings in a titre; therefore the total uncertainty in a titre is $2 \times 0.05 = 0.10 \text{ cm}^3$.

A reading involves temperature.
A measurement requires change (beginning temperature and end volume).



Occasionally, you might also have to factor in the uncertainty from judging an end point. There is an additional uncertainty of 0.05 cm^3 in every measurement.



Percentage uncertainties

Percentage uncertainties are a useful way of considering the scale of an uncertainty. 0.15 cm^3 (if including the uncertainty due to the drop) is not very big if the average titre is 17.4 cm^3 , but more important if the average titre is smaller, such as 8.5 cm^3 .

To calculate percentage uncertainty:

$$\% \text{ uncertainty} = \frac{\text{uncertainty}}{\text{value}} \times 100$$

Example

- a) Complete the results table.

	Rough	1	2	3
Start volume (cm^3)	0.00	18.15	0.05	17.35
End volume (cm^3)	18.60	36.25	17.40	34.80
Volume of acid (cm^3)				

	Rough
Start volume (cm^3)	0.00
End volume (cm^3)	18.60
Volume of acid (cm^3)	18.60

calculate

- b) Calculate the mean titre for the results to 2 decimal places.

$$\begin{aligned} \text{Mean titre} &= \frac{17.40 + 17.35}{2} \\ &= 17.425 \\ &= 17.43 \end{aligned}$$

- c) Calculate the percentage uncertainty of the mean titre due to the burette readings.

$$\begin{aligned} \% \text{ uncertainty} &= \frac{\text{uncertainty}}{\text{value}} \times 100 \\ &= \frac{0.1}{17.43} \times 100 \\ &= 0.574 \% \end{aligned}$$



COPYRIGHT
PROTECTED



Questions




Prepare your solutions

1. Complete the results table below for a titration experiment.

	Rough	1	2	3
Start volume (cm ³)	0.00	12.65	12.60	0.00
End volume (cm ³)	12.65	24.12	25.70	11.55
Volume of acid (cm ³)				

2. What are the readings on these burettes?



High concentration

3. A titration has the following results:

	Rough	1	2	3
Start volume (cm ³)	0.80	0.20	0.50	0.50
End volume (cm ³)	17.10	17.30	17.30	17.35
Volume of acid (cm ³)				

Find the mean titre for this titration.

4. Complete the results table below for a titration experiment, and calculate the mean titre.

	Rough	1	2	3
Start volume (cm ³)	0.05	19.50	0.10	19.05
End volume (cm ³)	19.50		19.05	38.05
Volume of acid (cm ³)		18.90		



Drop the base

5. The table below shows an incomplete set of results for a titration experiment. Calculate the mean titre, taking into account the uncertainty due to making readings (due to the volume of a drop).

	Rough	1	2	3
Start volume (cm ³)	0.00	0.00	0.40	0.15
End volume (cm ³)	22.55	43.10	43.05	43.05
Volume of acid (cm ³)				

6. In a titration experiment, a student fills the burette with acid. The burette was not cleaned before this use. Explain how this could affect the mean titre.

COPYRIGHT
PROTECTED



Concentrations and dilutions

Concentration

Concentrations are used in titrations to calculate the amounts of a substance in a solution. Concentration can be calculated using:

$$\text{concentration} = \frac{\text{moles}}{\text{volume}} \quad \text{and has units}$$

To work out how to make a solution with a specific concentration, you just have to know the concentration and the volume. For example, to make a solution with concentration 0.2 mol dm^{-3} and a volume of 300 cm^3 :



$$\text{moles} = \text{concentration} \times \text{volume} = 0.2 \times \frac{300}{1000} = 0.06 \text{ mol}$$

Converting units

In titrations you may be given a volume in cm^3 with a concentration in mol dm^{-3} . You need to convert from cm^3 to dm^3 .

$$1 \text{ dm}^3 = 1000 \text{ cm}^3 \text{ and } 1 \text{ cm}^3 = 0.001 \text{ dm}^3$$

To convert a volume from cm^3 to dm^3 , **divide** the value by 1000.

e.g. $20 \text{ cm}^3 = \frac{20}{1000} \text{ dm}^3 = 0.020 \text{ dm}^3$

To convert a volume from dm^3 to cm^3 , **multiply** the value by 1000.

e.g. $4.0 \text{ dm}^3 = 4.0 \times 1000 \text{ cm}^3 = 4000 \text{ cm}^3$





Diluting solutions

Diluting a solution by adding more solvent (e.g. distilled water) decreases its concentration. You can calculate the concentration of the diluted solution by dividing the concentration by the new volume.



$$\text{new concentration} = \frac{\text{old volume}}{\text{new volume}} \times \text{old concentration}$$

e.g. A solution with concentration 0.5 mol dm^{-3} is diluted with water from 5 cm^3 to 10 cm^3 .



$$\text{new concentration} = 0.5 \times \frac{5}{10} = 0.25 \text{ mol dm}^{-3}$$

For more complex dilutions, you may need to convert the concentrations into the same units before you perform the calculation, and then convert back into concentration at the end.



$$\text{concentration} = \frac{\text{moles}}{\text{volume}}$$

$$\text{concentration} =$$

COPYRIGHT
PROTECTED



Example 1

10 cm³ of a solution with concentration 0.30 mol dm⁻³ is diluted to 0.10 dm³. What is the concentration of the new solution?

$$\begin{aligned} 0.1 \text{ dm}^3 &= 100 \text{ cm}^3 \\ \text{Concentration} &= 0.30 \times \frac{10}{100} \\ &= 0.030 \text{ mol dm}^{-3} \end{aligned}$$

Example 2

Citric acid has an M_r of 192.1. 0.960 g of citric acid was dissolved in 50.0 cm³ of distilled water. The solution was then poured into a volumetric flask and made up to 250 cm³. Calculate the concentration of citric acid solution in the volumetric flask in mol dm⁻³.

$$\begin{aligned} \text{moles of citric acid} &= \frac{0.960}{192.1} \\ &= 0.00500 \\ \text{volume} &= \frac{250}{1000} \\ &= 0.250 \\ \text{concentration} &= \frac{0.00500}{0.250} \\ &= 0.0200 \text{ mol dm}^{-3} \end{aligned}$$

Questions



Prepare your solutions

- Find the concentration in g dm⁻³ of a solution where 3.00 g is dissolved in 500 cm³.
- How many moles are in 200 cm³ of a 0.200 mol dm⁻³ solution?
- Find the volume required to make a solution with concentration 0.500 mol dm⁻³ from 10.0 cm³ of a 2.00 mol dm⁻³ solution.



High concentration

- A standard solution of NaCl is made with concentration 2.00 mol dm⁻³. 25.0 cm³ of this solution is poured into a 250 cm³ volumetric flask. Distilled water is used to fill the flask to the mark. Calculate the concentration of NaCl in the new solution.
- Calculate the total volume of a solution of concentration 0.0125 mol dm⁻³ prepared by mixing 100 cm³ of a 0.400 mol dm⁻³ solution with water.
- In a titration, a 2.00 cm³ sample of Solution A is diluted to 100 cm³. The 100 cm³ sample is then used in a titration, and is found to have a concentration of 0.240 mol dm⁻³. Calculate the concentration of Solution A.



Drop the base

- A 200 cm³ solution of KI is made using 3.40 g of solid KI. KI has a relative formula mass of 166. Calculate the concentration of KI in mol dm⁻³.
- 250 cm³ of an NaCl solution with concentration 1.00 g dm⁻³ is diluted to give 1.00 dm³. Calculate the concentration of the new solution.

COPYRIGHT
PROTECTED



Structured acid-base titration calculations

Titration calculations

The purpose of a titration is to find the concentration of a solution. Acid-base titration involves a reaction between a solution of known concentration and a solution of unknown concentration.

You will need to use the equation: $\text{concentration} = \frac{\text{moles}}{\text{volume}}$

And to rearrange it to find: $\text{moles} = \text{concentration} \times \text{volume}$

and $\text{volume} = \frac{\text{moles}}{\text{concentration}}$

You will also need to consider the effects of ratios of reactants in the reaction equation.



ratio: 1 3 1 3

For this reaction, there is a 1 : 3 ratio of acid (H_3PO_4) to base (NaOH).


Finally, don't forget to convert your answer to the correct units.

Example 1

In a titration experiment, $0.210 \text{ mol dm}^{-3}$ NaOH was used to analyse a 25.0 cm^3 solution of HNO_3 .

	Rough	1	2	3
Start volume (cm^3)	0.00	0.00	0.00	0.00
End volume (cm^3)	45.80	45.65	45.50	45.35
Volume of acid (cm^3)	45.80			


The equation for this reaction is:



- Complete the results table above and calculate the mean titre.
- Calculate the number of moles of NaOH in the mean titre.
- Find the mass of HNO_3 dissolved in the 25.0 cm^3 solution to an appropriate number of significant figures. ($M_r \text{ HNO}_3 = 63.0$)

Example 2

In a titration experiment, a standard solution of H_2SO_4 was used to find the concentration of a solution of NaOH .



50.0 cm^3 samples of the NaOH solution were titrated against H_2SO_4 with a concentration of $0.200 \text{ mol dm}^{-3}$. The mean titre was 25.0 cm^3 .

- Find the number of moles of H_2SO_4 in the mean titre.
- Find the number of moles of NaOH in each sample.
- Find the concentration in mol dm^{-3} of the NaOH solution to three significant figures.

a)

Start volume (cm^3)
End volume (cm^3)
Volume of acid (cm^3)

mean titre = $\frac{45.65 + 45.50}{2} = 45.575$

b) $\text{mol NaOH} = \frac{45.575}{1000} \times 0.210 = 0.00957$

c) ratio in reaction 1 : 1
 $\text{mol HNO}_3 = 0.00957$

mass $\text{HNO}_3 = 0.00957 \times 63.0 = 0.603 \text{ g}$

$\text{mol} = \text{conc} \times \text{vol}$

a) $\text{mol H}_2\text{SO}_4 = \frac{25.0}{1000} \times 0.200 = 0.005$

b) Ratio in reaction 1 : 2

$\text{mol NaOH} = 2 \times 0.005 = 0.01$

c) $\text{conc NaOH} = \frac{0.01}{0.05} = 0.200$

INSPECTION COPY

COPYRIGHT
PROTECTED



Questions




Prepare your solutions

1. In a titration experiment, $0.200 \text{ mol dm}^{-3}$ HCl is put into a burette. The following results are recorded:

	Rough	1	2	3
Start volume (cm^3)	0.00	1.10	0.10	16.15
End volume (cm^3)	16.25	2.25	16.15	32.25
Volume of acid (cm^3)				

- Find the mean titre for this titration.
 - Calculate the moles of HCl used in the mean titre.
2. In a titration experiment, a solution of H_2SO_4 with a concentration of $0.100 \text{ mol dm}^{-3}$ is added to a solution of NaOH.



The following results are recorded:

	Rough	1	2	3
Start volume (cm^3)	0.00	0.00	23.85	0.00
End volume (cm^3)	24.40	23.85	47.65	23.80
Volume of acid (cm^3)				

- Find the mean titre for this titration.
- Calculate the moles of H_2SO_4 used in the mean titre.
- Calculate the number of moles of NaOH which reacted with the H_2SO_4 .



High concentration

3. A student performs a titration experiment between HCl and KOH.

30.0 cm^3 samples of KOH were titrated with $0.200 \text{ mol dm}^{-3}$ HCl. The student recorded a mean titre of 13.70 cm^3 .

- Calculate the moles of HCl in the mean titre.
 - Calculate the moles of KOH in the 30.0 cm^3 samples.
 - Calculate the concentration of KOH.
4. A student performs a titration experiment between H_2SO_4 and KOH.

80.0 cm^3 samples of KOH were titrated with H_2SO_4 . The H_2SO_4 solution was prepared by diluting a 2.00 mol dm^{-3} solution to 1.00 dm^3 . The student calculated the mean titre to be 16.00 cm^3 .

- Calculate the concentration of the H_2SO_4 used in the titration.
- Calculate the moles of H_2SO_4 in the mean titre.
- Calculate the moles of KOH in the 80.0 cm^3 samples.
- Calculate the concentration of KOH.

**COPYRIGHT
PROTECTED**






Drop the base

5. A company wants to use calcium hydroxide to pickle cucumbers. To do this, it needs a solution of calcium hydroxide that has a concentration of at least 6.00 g dm^{-3} .

A scientist diluted a solution of Ca(OH)_2 from 200 cm^3 to 1 dm^3 . The scientist then used $0.206 \text{ mol dm}^{-3}$ HCl to titrate 250 cm^3 samples of the solution of Ca(OH)_2 . The following results were produced:

	Rough	1	2	3
Start volume (cm^3)	0.05	0.10	0.40	0.20
End volume (cm^3)	1.05	40.55	40.50	40.50
Volume of acid (cm^3)				

- Use the results table to find the mean titre.
 - Calculate the moles of Ca(OH)_2 in the 250 cm^3 samples to a suitable number of significant figures.
 - Calculate the concentration of Ca(OH)_2 in the original 200 cm^3 solution in g dm^{-3} .
 - Determine whether this sample of calcium hydroxide is concentrated enough for pickling cucumbers.
6. A titration experiment involves the following reaction of a base, MOH, with HCl:



236.4 mg of MOH was dissolved in 200.0 cm^3 of water. 20.0 cm^3 samples of the solution were taken, and 100 cm^3 of $0.245 \text{ mol dm}^{-3}$ HCl. The following results table was produced:

	Rough	1	2	3
Start volume (cm^3)	0.55	0.55	0.55	0.40
End volume (cm^3)	18.15	17.55	17.70	17.65
Volume of acid (cm^3)				

- Calculate the mean titre and determine the number of moles of HCl in the mean titre.
- Calculate the number of moles of MOH in the 20.0 cm^3 samples, and then determine the concentration of MOH to an appropriate number of significant figures.

INSPECTION COPY

COPYRIGHT
PROTECTED



Unstructured acid-base titration calculations

Titration Calculations

Titration calculations can be used to find out a number of different things, such as:

- the concentration of an unknown solution
- the mass of compound in an unknown substance
- the M_r of a compound / the identity of a compound from a list using M_r


The steps that lead up to answers to these questions are usually very similar, and you will need to be able to carry them out comfortably under exam conditions. The steps are labelled on the two examples below.

1. Work out the **equation** for the reaction that happens in the titration.
2. Work out the **moles** in the mean titre.
3. Use the mole **ratio** to work out the moles in the solution being titrated.
4. Work out the moles in the **original** solution, i.e. before it was diluted or samples were taken for titration.
5. **Answer** the question.

Example 1

In a titration experiment, a standard solution of HNO_3 was used to find the concentration of a solution of KOH .

Step 1 (given in question)



20.0 cm^3 samples of the NaOH solution were titrated against H_2SO_4 with a concentration of 0.150 mol dm^{-3} . The mean titre was 18.95 cm^3 .

Find the concentration of the NaOH solution to three significant figures.

Step 2

$$\begin{aligned} \text{mol HNO}_3 &= 0.150 \times \frac{18.95}{1000} \\ &= 0.0028425 \text{ mol} \end{aligned}$$

Step 3

from reaction $\text{HNO}_3 : \text{NaOH}$

Step 4

$$\text{mol KOH} = 0.0028425$$

Step 5

$$\begin{aligned} \text{conc KOH} &= 0.0028425 \div 0.0200 \\ &= 0.142125 \text{ mol dm}^{-3} \\ &\quad \swarrow \\ &\quad 3 \text{ s.f.} \end{aligned}$$

Example 2

Compound A is one of the monoprotic carboxylic acids from the list below:

- methanoic acid HCOOH
- ethanoic acid CH_3COOH
- propanoic acid $\text{CH}_3\text{CH}_2\text{COOH}$
- butanoic acid $\text{CH}_3\text{CH}_2\text{CH}_2\text{COOH}$

5760 mg of Compound A is mixed with water and made up to 500.0 cm^3 . 50.0 cm^3 samples of the solution are titrated against 0.200 mol dm^{-3} NaOH .

The mean titre of NaOH is 5.76 cm^3 . Use the results to identify Compound A.

Step 2

$$\begin{aligned} \text{moles NaOH} &= 0.200 \times \frac{5.76}{1000} \\ &= 0.00655 \text{ mol} \end{aligned}$$

Step 3

monoprotic, so 1 : 1 ratio

$$n(\text{A}) \text{ in } 25 \text{ cm}^3 = 0.00655 \text{ mol}$$

$$\begin{aligned} n(\text{A}) \text{ in } 500 \text{ cm}^3 &= 0.00655 \times 20 \\ &= 0.0655 \text{ mol} \end{aligned}$$

Step 4

$$\begin{aligned} M_r \text{ A} &= \frac{5760}{0.0655} \\ &= 87.9 \end{aligned}$$

Step 5

Compound A is butanoic acid

INSPECTION COPY

COPYRIGHT
PROTECTED



Questions




Prepare your solutions

- 50.0 cm³ samples of NaOH with unknown concentration were titrated against 0.100 mol dm⁻³ HCl, giving the following results. Calculate the concentration of NaOH.



	Rough	1	2	3
Start volume (cm ³)	0.00	18.70	0.65	17.65
End volume (cm ³)	20.70	35.45	17.65	34.70
Volume of acid (cm ³)				

- 25.0 cm³ samples of KOH with concentration 0.130 mol dm⁻³ were titrated with 0.250 mol dm⁻³ H₂SO₄, giving the following results. Calculate the concentration of H₂SO₄.




	Rough	1	2	3
Start volume (cm ³)	0.45	0.55	0.00	0.40
End volume (cm ³)	32.55	31.50	31.55	31.35
Volume of acid (cm ³)				



High concentration

- 1.37 g of an unknown hydroxide MOH is dissolved in water and made up to 250 cm³. 25.0 cm³ of this solution is titrated against 0.100 mol dm⁻³ H₂SO₄, giving a mean titre of 30.58 cm³. Use this to calculate the relative formula mass of the hydroxide MOH.



- The purity of a solid sample of K₂CO₃ is tested by titration. 9.33 g of the sample is dissolved in water and made up to 250 cm³. 25.0 cm³ of this solution are titrated with 0.125 mol dm⁻³ HCl.

	Rough	1	2	3
Start volume (cm ³)	0.55	0.30	0.05	0.15
End volume (cm ³)	41.90	41.10	41.10	41.05
Volume of acid (cm ³)				


Write a balanced equation for the reaction between HCl and K₂CO₃, which produces two products, and use the results to find the purity of the solid sample.

$$\text{percentage purity} = \frac{\text{mass of K}_2\text{CO}_3}{\text{mass of sample}} \times 100$$



Drop the base

- Two 10 cm³ solutions of KOH, Solution A and Solution B, are mixed together in a volumetric flask. Solution A has a concentration of 0.100 mol dm⁻³, and Solution B has a concentration of 0.200 mol dm⁻³. 25.0 cm³ samples of the mixture are titrated with 0.200 mol dm⁻³ H₂SO₄, giving a mean titre of 21.80 cm³. Write the symbol equation for this reaction, and use it to calculate the concentration of Solution B.
- A 5.34 g sample contains 2.86 g of Ba(OH)₂, an unknown mass of NaOH, and an impurity. The sample is dissolved in water and made up to 250 cm³ in a volumetric flask. 25.0 cm³ of this solution are titrated with 0.150 mol dm⁻³ H₂SO₄, giving a mean titre of 28.85 cm³.



Find the mass of the unreactive impurity in the sample in g.

COPYRIGHT
PROTECTED



'Back' titrations

'Back' titration


A back titration is a type of titration used to investigate substances which are not very soluble, or when a reaction would be too slow to study using titration. For example, to work out the amount of calcium carbonate in a rock, we cannot use a normal titration because calcium carbonate does not dissolve in water. Instead we would react the calcium carbonate with an excess of acid first, in a back titration.

There are normally two steps to the back titration:

1. An **excess** of acid or alkali is added so that all of the compound being analysed reacts.
2. The amount of acid/alkali remaining is then found using titration.

The amount of titrant is calculated by:

- finding out how much acid/alkali is remaining and using this value to calculate the amount of the original substance
- using how much acid/alkali was used up to work out how much of the unknown original solution to react with it




Example 1

A 132 mg solid sample of a carbonate (MCO_3), is added to 50.0 cm^3 of HCl of concentration $0.120 \text{ mol dm}^{-3}$. The reaction is:



The resulting solution is analysed with $0.100 \text{ mol dm}^{-3} \text{ NaOH}$, giving a mean titre of 28.68 cm^3 .



Determine the identity of the carbonate.

Step 1: find the moles of NaOH

$$\begin{aligned} \text{mol NaOH} &= 0.100 \times \frac{28.68}{1000} \\ &= 0.002868 \text{ mol} \end{aligned}$$

$$\text{mol HCl} = 0.002868$$

Step 2: calculate the moles of HCl

$$\begin{aligned} \text{total mol HCl} &= 0.120 \times \frac{50}{1000} \\ &= 0.00600 \end{aligned}$$

$$\begin{aligned} \text{mol HCl used} &= 0.00600 - 0.002868 \\ &= 0.003132 \text{ mol} \end{aligned}$$

Step 3: calculate the moles of MCO_3

$$\begin{aligned} \text{mol MCO}_3 &= \frac{0.003132}{2} \\ &= 0.001566 \text{ mol} \end{aligned}$$

Step 4: answer the question

$$\begin{aligned} M_r \text{ of MCO}_3 &= \frac{132}{0.001566} \\ &= 84.3 \\ A_r \text{ of M} &= 84.3 - 12 \\ \text{MCO}_3 &\text{ is MgCO}_3 \end{aligned}$$

**COPYRIGHT
PROTECTED**



Questions




Prepare your solutions

1. A 0.782 g sample of chalk contains CaCO_3 and some unreactive impurities. The solution of $0.100 \text{ mol dm}^{-3}$ HCl.



50.0 cm^3 samples of the resulting solution are titrated with $0.100 \text{ mol dm}^{-3}$ KOH. 17.88 cm^3 .



- Calculate the moles of KOH in the mean titre.
 - Calculate the moles of HCl remaining in the 250 cm^3 solution after the CaCO_3 reaction.
 - Calculate the moles of HCl which reacted with the CaCO_3 .
 - Calculate the moles of CaCO_3 added to the acid.
 - Calculate the mass of calcium carbonate present in the rock sample.
2. Ammonia is a volatile base and cannot be titrated against directly. A back titration is used to find the number of moles of ammonia in a 3.00 dm^3 solution.

A 30.0 cm^3 sample of the 3.00 dm^3 solution reacts with 0.0250 mol of HCl and distilled water.



20.0 cm^3 samples of the resulting solution are titrated against $0.100 \text{ mol dm}^{-3}$ KOH. 32.45 cm^3 .

- Calculate the moles of KOH in the mean titre.
- From the amount of KOH reacted, calculate the moles of HCl remaining in the 20.0 cm^3 sample after reaction with NH_3 .
- Calculate the moles of HCl which reacted with the NH_3 .
- Calculate the moles of ammonia in the 3.00 dm^3 solution.

INSPECTION COPY


COPYRIGHT
PROTECTED





High concentration

3. 1.15 g of a pure solid sample of a hydroxide with the formula $M(OH)_2$ is placed in 25.0 cm³ of 0.100 mol dm⁻³ H_3PO_4 and diluted to 250 cm³.




25.0 cm³ samples of the resulting solution are titrated against KOH with concentrated KOH solution.



The following results are recorded.

	Rough	1	2	3
Start volume (cm ³)	0.55	0.15	0.05	0.20
End volume (cm ³)	40.00	39.00	38.95	39.00
Volume of acid (cm ³)				



- Balance the equation for the reaction between $M(OH)_2$ and H_3PO_4 .
 - From the mean titre, find the amount in moles of H_3PO_4 in 250 cm³ after reaction.
 - Determine the formula of the salt $M(OH)_2$.
4. A pure solid is thought to be sodium carbonate heptahydrate, $Na_2CO_3 \cdot 7H_2O$. Determine the relative formula mass of the salt by reacting the solid sodium carbonate with a standard solution of 0.100 mol dm⁻³ HNO_3 and then titrating the resulting solution with a standard solution of 0.100 mol dm⁻³ $NaOH$. Outline, with brief practical details, an experiment to determine the relative formula mass of the solid, including any calculations you would perform. You are provided with 1.00 g of the solid.



Drop the acid


5. An insoluble solid salt has the formula $MgCO_3 \cdot xH_2O$, where x is a whole number. Determine the relative formula mass of the salt. A crucible is placed on a balance and weighed. The mass is 142.34 g. The salt is placed in the crucible, and the combined mass of the crucible and salt is 157.34 g. This mass of $MgCO_3 \cdot xH_2O$ is placed in 150.0 cm³ of 0.240 mol dm⁻³ HCl and reacted. The solution is diluted to 250 cm³, and 50.0 cm³ samples are titrated with 0.100 mol dm⁻³ $NaOH$ giving a mean titre of 20.08 cm³. Calculate the value of x .

6. A 200 cm³ solution contains a mixture of $Ba(OH)_2$ and $NaOH$. 250.0 cm³ H_2SO_4 of 0.200 mol dm⁻³ is added so that H_2SO_4 is in excess of both reagents. The following reactions occur:



$BaSO_4$ is a precipitate and is removed by filtration. The filtrate is dried and weighed to be 1.00 g.

The remaining solution is made up to 500 cm³, and 50.0 cm³ samples are titrated with 0.100 mol dm⁻³ HCl giving a mean titre of 41.30 cm³.



Calculate the concentration of $NaOH$ in the 200 cm³ solution in mol dm⁻³.

COPYRIGHT
PROTECTED




Titration curves / indicators

Titration curves

Titration curves show how pH changes as one reagent is added to the other in a reaction. The graph shows a simple neutralisation curve for a titration in which $0.100 \text{ mol dm}^{-3}$ NaOH is added to 25 cm^3 of $0.100 \text{ mol dm}^{-3}$ HCl.

Key points

- The graph represents the change in pH as NaOH is added to HCl.
- The solution starts acidic (low pH) and becomes more alkaline (high pH).
- At the equivalence point, a small volume of NaOH added leads to a large change in pH. This is where the indicator changes colour.
- The equivalence point is the centre of the steep vertical section of the graph.
- For this titration, the acid and base react in a 1 : 1 ratio, and have equal concentration, so the equivalence point occurs when the volumes are equal (25 cm^3).





Strong and weak acids and bases

strong acid



weak acid

strong base



Choice of indicator depends on the pH at the equivalence point of the neutralisation reaction.

weak base



COPYRIGHT
PROTECTED





INSPECTION COPY

Example 1

Choose a suitable indicator from the list below for the following graph.

Indicator	pH range
methyl orange	3.1–4.4
bromothymol blue	6.0–7.6
phenolphthalein	8.3–10.0




Step section is from approx

Methyl orange is the only su

Example 2

Draw a pH curve for 0.200 mol dm⁻³ NaOH, a strong base, added to 30 cm³ 0.100 mol dm⁻³ CH₃COOH, a weak acid.


Label the equivalence point.



Example 3

Draw a pH curve for 0.100 mol dm⁻³ HCl, a strong acid, added to 20 cm³ 0.100 mol dm⁻³ NaOH, a strong base.

Label the equivalence point.



COPYRIGHT
PROTECTED




Questions



Prepare your solutions

Here are four titration curves. All reagents have a concentration of $0.100 \text{ mol dm}^{-3}$



- Write the letter of the curve produced by
 - adding hydrobromic acid to 25 cm^3 potassium hydroxide
 - adding ammonia to 25 cm^3 propanoic acid
 - adding sodium hydroxide to 25 cm^3 ethanoic acid
- This table shows information about some pH indicators which could be used

Indicator	pH range	Low pH colour
bromothymol blue	6.0–7.6	yellow
phenolphthalein	8.3–10.0	colourless
alizarine yellow	10.1–12.0	yellow
bromocresol green	3.8–5.4	yellow
thymolphthalein	9.3–10.5	colourless

- Which indicator(s) could be used to track the titration represented by curve A?
- Which indicator could be used to track the titration represented by curve B?
- Identify the colour change for thymolphthalein during the titration represented by curve C.



High concentration

- A student has a solution with a pH meter and finds it has a pH of 8.2. The student adds phenolphthalein, bromocresol green and thymolphthalein. Assume that none of the indicators reacts with the solution and that the pH of the solution does not change. Use the table above to identify which indicator(s) would be most suitable for the titration.
- Sketch a titration curve for 50 cm^3 $0.100 \text{ mol dm}^{-3}$ HCl added to 35 cm^3 $0.100 \text{ mol dm}^{-3}$ NaOH.

INSPECTION COPY

COPYRIGHT
PROTECTED





Drop the base

5. Sketch the titration curve for 50 cm^3 $0.100 \text{ mol dm}^{-3}$ NH_3 added to 10 cm^3 0.20 mol dm^{-3} HCl .
6. A student is performing a series of titrations of the acids and bases shown in the table below.

Acid		Base	
weak	strong	weak	strong
CH_3COOH	HCl	CH_3NH_2	KOH

The following indicators are available:



Indicator	pH range
methyl orange	3.1–4.4
bromothymol blue	6.0–7.6
phenolphthalein	8.3–10.0

For the following titrations, select the indicators which would give a precise end point.

- CH_3COOH titrated against KOH
- CH_3NH_2 titrated against HBr
- HBr titrated against KOH



INSPECTION COPY

COPYRIGHT
PROTECTED



Strong acids and bases

Brønsted-Lowry

According to the Brønsted-Lowry theory of acids and bases:

Acid	Base
proton donor	proton acceptor
$HA \rightleftharpoons H^+ + A^-$	$B + H_2O \rightleftharpoons BH^+ + OH^-$

Strong acids

Strong acids are acids which completely ionise in solution. This means we can write **irreversibly**. The concentration of H^+ ions can be found from the concentration of


- HCl releases one H^+ ion, so $[H^+] = [HCl]$ $HCl \rightarrow H^+ + Cl^-$
- H_2SO_4 releases two H^+ ions, so $[H^+] = 2 \times [H_2SO_4]$ $H_2SO_4 \rightarrow 2H^+ + SO_4^{2-}$

pH

pH is a way of measuring the concentration of H^+ ions.

You can find $[H^+]$ from pH using:

$$[H^+] = 10^{-pH}$$



Strong bases

Bases are in equilibrium: $HA + H_2O \rightleftharpoons BH^+ + OH^-$
 To calculate $[H^+]$ for a base, first find $[OH^-]$.
 To do this you have to use K_w , the ionic product of water.

At room temperature:

$$K_w = [H^+] \times [OH^-] = 1.00 \times 10^{-14}$$

To calculate $[H^+]$, rearrange this equation: $[H^+] = \frac{1.00 \times 10^{-14}}{[OH^-]}$

The pH can be calculated in the usual way: $pH = -\log_{10}[H^+]$



Example 1

Find the pH of a solution of H_3PO_4 with concentration $0.00120 \text{ mol dm}^{-3}$.

$$\begin{aligned} [H^+] &= 3 \times 0.00120 \\ &= 0.00360 \text{ mol dm}^{-3} \\ pH &= -\log_{10}(0.00360) \\ &= 2.44 \end{aligned}$$

Example 2

Find the concentration of a solution of HCl with pH 1.3.

$$\begin{aligned} \text{conc} &= 10^{-1.3} \\ &= 0.0501 \end{aligned}$$

Example 3

Find the pH of a solution of $NaOH$ with concentration $0.00670 \text{ mol dm}^{-3}$ at room temperature. ($K_w = 1.0 \times 10^{-14}$)

$$\begin{aligned} [OH^-] &= 0.00670 \text{ mol dm}^{-3} \\ [H^+] &= \frac{1.00 \times 10^{-14}}{0.00670} \\ &= 1.49 \times 10^{-12} \\ pH &= -\log_{10}(1.49 \times 10^{-12}) \\ &= 11.8 \end{aligned}$$

INSPECTION COPY

COPYRIGHT
PROTECTED



Questions




Prepare your solutions

- For the following solutions, convert between pH and concentration.
 - HCl concentration $0.00100 \text{ mol dm}^{-3}$
 - HBr pH 2.30
 - H_2SO_4 concentration $0.000400 \text{ mol dm}^{-3}$
- Find the pH of the following strong bases using the ionic product of water K_w .
 - NaOH concentration $0.00160 \text{ mol dm}^{-3}$
 - KOH concentration $0.000810 \text{ mol dm}^{-3}$
 - Ca(OH)_2 concentration $0.0000451 \text{ mol dm}^{-3}$



High concentration

- Ammonia, NH_3 , is added to water to form a solution for use in a titration. During this step, the following equilibrium occurs:



The pH of the solution for titrating is found to be 11.7.


Calculate the concentration of OH^- ions in the solution using the ionic product $K_w = 1.0 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$.

- A solution of H_2SO_4 is found by titration to have a concentration of $0.000520 \text{ mol dm}^{-3}$. Calculate the concentration of OH^- ions in the solution.



Drop calculations

- In a titration experiment, 500 cm^3 of $0.0100 \text{ mol dm}^{-3}$ HCl is added to 150 cm^3 of $0.0200 \text{ mol dm}^{-3}$ NaOH. In this step, the following reaction occurs:



By working out the concentration of either the H^+ ions or the OH^- ions left over, calculate the pH of the resulting solution.

- At 10°C , the ionic product of water is 2.88×10^{-15} .
 - Use a calculation to show whether pH 7 is acidic, neutral or alkaline at 10°C .
 - 40.0 cm^3 $0.00420 \text{ mol dm}^{-3}$ KOH is diluted to 800 cm^3 at 10°C . Calculate the pH of the resulting solution at 10°C to 2 d.p.

INSPECTION COPY

COPYRIGHT
PROTECTED



Weak acids

Weak acids

Acids are substances that dissociate to form H^+ ions. Whereas strong acids dissociate completely. Many organic acids, e.g. carboxylic acids, are weak acids.



K_a

Each acid has a quantity called its K_a which is a way of measuring the strength of an acid. The more it dissociates, the higher the K_a value.

For an acid $HA \rightleftharpoons H^+ + A^-$

K_a can be calculated as:

When the acid HA dissociates, equal numbers of A^- and H^+ ions are in the solution.

$$[A^-] = [H^+]$$

and so:

$$K_a = \frac{[H^+]^2}{[HA]} \quad \text{which can be used to calculate } K_a$$

The stronger an acid, the more of it dissociates to form H^+ , and, therefore, the higher the K_a value.

pK_a

K_a values are often extremely small or extremely large numbers, so K_a values are converted to **pK_a values** to make them more manageable.

pK_a is calculated as:

$$pK_a = -\log_{10}(K_a)$$

Stronger acids have lower (or even negative) pK_a values.


Example 1

Phenol has a pK_a value of 9.80. Calculate the K_a of phenol.

$$\begin{aligned} K_a &= 10^{-pK_a} \\ &= 10^{-9.8} \\ &= 1.6 \times 10^{-10} \end{aligned}$$

Example 2

CCl_3COOH has a pK_a value of 0.65. Find the concentration of H^+ ions in a $0.0500 \text{ mol dm}^{-3}$ solution of CCl_3COOH .



$$\begin{aligned} K_a &= 10^{-pK_a} \\ &= 10^{-0.65} \\ &= 0.224 \end{aligned}$$

$$K_a = \frac{[CCl_3COO^-][H^+]}{[CCl_3COOH]}$$


$$[H^+] = [CCl_3COO^-], \text{ so}$$

$$K_a = \frac{[H^+]^2}{[CCl_3COOH]}$$

$$\begin{aligned} [H^+] &= \sqrt{[CCl_3COOH] \times K_a} \\ &= \sqrt{0.05 \times 0.224} \\ &= 0.110 \text{ mol dm}^{-3} \end{aligned}$$

Example 2

CH_3COOH has a K_a value of 1.58×10^{-5} .



Calculate the pH of a $0.200 \text{ mol dm}^{-3}$ solution of CH_3COOH .

$$K_a = \frac{[CH_3COO^-][H^+]}{[CH_3COOH]}$$

$$[H^+] = [CH_3COO^-], \text{ so}$$

$$K_a = \frac{[H^+]^2}{[CH_3COOH]}$$

$$\begin{aligned} [H^+] &= \sqrt{[CH_3COOH] \times K_a} \\ &= \sqrt{0.2 \times 1.58 \times 10^{-5}} \\ &= 1.78 \text{ mol dm}^{-3} \end{aligned}$$

$$\begin{aligned} \text{pH} &= -\log_{10}(1.78) \\ &= -0.25 \end{aligned}$$

INSPECTION COPY

COPYRIGHT
PROTECTED



Questions



Prepare your solutions

- Convert these values:
 - $K_a = 1.42 \times 10^{-5}$ to pK_a
 - $K_a = 4.98 \times 10^{-7}$ to pK_a
 - $pK_a = 1.03$ to K_a
 - $pK_a = 3.12$ to K_a
- HSCN is a weak acid with a pK_a of 1.1.
 - Write an expression for the K_a of HSCN.
 - Calculate the pH of $0.0108 \text{ mol dm}^{-3}$ HSCN.



High concentration

- HNO_2 is a weak acid with a pK_a of 3.29.
A solution of HNO_2 is found to have a concentration of $0.0452 \text{ mol dm}^{-3}$.
Calculate the concentration of NO_2^- ions in a solution.
- A $0.0230 \text{ mol dm}^{-3}$ solution of phenol, $\text{C}_6\text{H}_5\text{OH}$, has a pH of 5.79. Calculate the



Drop the base

- pK_a values are often given for acids dissolved in water. pK_a can also be given for acids dissolved in other solvents, for example DMSO.

The table below shows the pK_a values of ethanoic acid, CH_3COOH , in two solvents.

pK_a in water	pK_a in DMSO
4.76	12.3

- Explain what it means that ethanoic acid has a higher pK_a in DMSO than in water.
 - What concentration of ethanoic acid in water would have the same pH as ethanoic acid dissolved in DMSO?
- The pK_a of water, H_2O , can be calculated in the same way as that of any other acid.
 - Write an expression for the K_a of water.
 - 1 dm^3 of H_2O contains 1000 g of water. Determine the value of $[\text{H}_2\text{O}]$ given the formula mass of 18.
 - Calculate the pK_a of water using the concentration of H_2O along with $[\text{H}^+]$ and the product of water $[\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14}$.

**COPYRIGHT
PROTECTED**



Buffers


Buffers

Buffers are solutions which can **minimise** the change in pH when a **small amount**

Forming buffers


Buffers are formed by combining a weak acid with a salt of the weak acid. For example, a buffer could be formed by adding sodium ethanoate to ethanoic acid.

Sodium ethanoate salt dissociates in solution:
Ethanoic acid is in equilibrium:



How buffers minimise pH changes

Buffers minimise pH changes when H^+ or OH^- ions are added through shifts in the base equilibrium:



For the examples of $\text{CH}_3\text{COOH}/\text{CH}_3\text{COO}^-$ buffer above:

When H^+ ions are added

- The concentration of H^+ increases.
- The equilibrium $\text{CH}_3\text{COOH} \rightleftharpoons \text{CH}_3\text{COO}^- + \text{H}^+$ shifts to the left and $[\text{H}^+]$ goes back down.
- The pH change is minimised.

When OH^- ions are added

- OH^- reacts with H^+ .
- The concentration of H^+ decreases.
- The position of equilibrium shifts to the right and $[\text{H}^+]$ goes back up.
- The pH change is minimised.

Buffer calculations

Buffer calculations often involve finding the pH of a buffer solution. It is important to rearrange the formula correctly.

For this, use the formula:

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

Which rearranges to:

$$[\text{H}^+] = K_a \times \frac{[\text{HA}]}{[\text{A}^-]}$$

Once you have calculated $[\text{H}^+]$, you can calculate pH in the usual way, using $-\log_{10}[\text{H}^+]$.

Example 1

A 1 dm³ buffer solution is formed by mixing 400 cm³ 0.175 mol dm⁻³ CH₂ClCOOH (acid) with 600 cm³ 0.200 mol dm⁻³ CH₂ClCOONa (salt).

CH₂ClCOOH has a K_a of 1.380×10^{-3} . Calculate the pH of the buffer solution.

First, you need to calculate the concentration of the acid and the salt when they have been mixed.

$$\begin{aligned} [\text{HA}] &= \frac{0.175 \times 400}{1000} \\ &= 0.0700 \\ [\text{A}^-] &= \frac{0.200 \times 600}{1000} \\ &= 0.120 \\ [\text{H}^+] &= 1.380 \times 10^{-3} \times \frac{0.0700}{0.120} \\ &= 8.05 \times 10^{-4} \text{ mol dm}^{-3} \\ \text{pH} &= 3.09 \end{aligned}$$

INSPECTION COPY

COPYRIGHT
PROTECTED



Example 2

A scientist wants to make a buffer with a pH of 3.60. Calculate the mass of NaNO_2 which could be added to a 150 cm^3 $0.200 \text{ mol dm}^{-3}$ solution of HNO_2 . HNO_2 has a pK_a of 3.29.

The first step is to recognise from the NaNO_2 is A^- .

$$\begin{aligned} [\text{A}^-] &= K_a \times \frac{[\text{HA}]}{[\text{H}^+]} \\ K_a &= 10^{-3.29} \\ &= 5.129 \times 10^{-4} \\ [\text{H}_2\text{O}] &= 0.200 \text{ mol dm}^{-3} \\ [\text{H}^+] &= 10^{-3.6} \\ &= 2.512 \times 10^{-4} \text{ mol dm}^{-3} \\ [\text{A}^-] &= 5.129 \times 10^{-4} \times \frac{0.200}{2.512 \times 10^{-4}} \\ &= 0.408 \text{ mol dm}^{-3} \\ \text{mol A}^- &= 0.408 \times \frac{150}{1000} \\ &= 0.0613 \text{ mol} \\ \text{mass HNO}_2 &= 0.0613 \times 69 \\ &= 4.22 \text{ g} \end{aligned}$$

$$\text{mol HNO}_2 = \text{mol A}^-$$

Questions



Prepare your solutions

- Suggest a salt that could be used to form a buffer with the following weak acids
 - Methanoic acid, CHOOH
 - HCN
- A buffer solution is created by mixing a solution of CF_3COOH with CF_3COONa . the buffer equilibrium is shown. This buffer system minimises changes in pH when
 - a small amount of HCl is added
 - a small amount of NaOH is added



High concentration

- A 1.00 dm^3 buffer solution is formed by mixing 200 cm^3 of $0.150 \text{ mol dm}^{-3}$ H_2S with $0.150 \text{ mol dm}^{-3}$ HS^- .

 H_2S has a pK_a of 7.04. Calculate the pH of the buffer solution.
- A small amount of ammonia, NH_3 , is added to an excess of hydrofluoric acid, HF , to form a buffer which regulates pH when an acid or an alkali is added.
- Lactic acid, $\text{CH}_3\text{CH}(\text{OH})\text{COOH}$, has a pK_a of 3.86. A mass of lactic acid is added to a solution of $\text{NaOOCCH}_2\text{CH}_3$.
 - Explain why a buffer forms.
 - Find the mass of lactic acid needed to create a buffer solution with a pH of 3.86.



Drop the base

- A buffer is formed between 600 cm^3 of $0.350 \text{ mol dm}^{-3}$ $\text{C}_5\text{H}_4\text{NHCl}$ pK_a 5.25 and a solution containing its conjugate base. Find the mass of the conjugate base required to create a buffer system to 5.60.

COPYRIGHT
PROTECTED



Structured redox titration

Redox

Redox reactions occur when one species' oxidation number (also known as oxidation state) increases and another species' oxidation number decreases.

This worksheet is for understanding oxidation numbers.

An increase in oxidation number is called **oxidation**. A species which causes oxidation is called an **oxidising agent**. A species which has a reduction in oxidation number is **reduced**. Something that causes reduction is called a **reducing agent**.

e.g. $2\text{Fe}^{3+} + \text{Cu} \rightarrow 2\text{Fe}^{2+} + \text{Cu}^{2+}$

Fe^{3+} goes from $3+ \rightarrow 2+$ so is reduced and is, therefore, an oxidising agent.

Cu goes from $0 \rightarrow 2+$ so is oxidised and is, therefore, a reducing agent.


Redox titrations

Redox titrations use redox reactions rather than acid-base neutralisations. Because they involve a colour change, many redox titrations don't need an indicator.


Whereas equations for neutralisation reactions used in acid-base titrations are usually simple, redox reactions can be quite complex.

Balancing redox equations

Firstly, the individual half-equations must be balanced, and then the equations combined. e.g.



This equation is balanced.




This equation is not balanced.


- **Step 1: Balance the atoms.** – use $\text{H}^+/\text{H}_2\text{O}$ (acid) or $\text{OH}^-/\text{H}_2\text{O}$ if you need to.




- **Step 2: Balance the charges.**



- **Step 3: Combine the two equations to make a balanced redox equation.**
Combine the reactions so that the electrons balance on each side:



Cancel the electrons (sometimes you will have to cancel other species such as H^+ or OH^-).




INSPECTION COPY

COPYRIGHT
PROTECTED



Example 1

Iodine (I_2) and thiosulfate ($S_2O_3^{2-}$) react together in the following reaction:




50.0 cm³ samples of an iodine solution were titrated with 0.120 mol dm⁻³ thiosulfate solution. The mean titre was 26.78 cm³.

- Calculate the number of moles of thiosulfate used in the mean titre.
- Calculate the number of moles of iodine required to react with this amount of thiosulfate.
- Calculate the concentration of iodine in the original solution.

$$\begin{aligned} \text{a) } \text{mol } S_2O_3^{2-} &= \frac{26.78}{1000} \times 0.120 \\ &= 0.00535 \\ \text{b) } \text{mol } I_2 &= 0.00535 \div 2 \\ &= 0.002675 \\ \text{c) } \text{conc } I_2 &= 0.002675 \div 0.050 \\ &= 0.0535 \text{ mol dm}^{-3} \end{aligned}$$

Example 2

A lump of soluble iron(II) salt was dissolved in 250 cm³ water, and 25.0 cm³ samples were taken and titrated with 0.150 mol dm⁻³ MnO_4^- solution. The titres recorded were 34.55, 34.50 and 34.70 cm³.



- Use the concordant titres to calculate the mean volume of MnO_4^- in cm³.
- Use the mean titre to calculate the moles of MnO_4^- in the titration.
- Calculate the moles of Fe^{2+} titrated.
- Calculate the moles of Fe^{2+} in the original lump of salt.
- Calculate the total mass of iron in the original Fe^{2+} salt.

$$\begin{aligned} \text{a) } \text{vol } MnO_4^- &= \frac{34.55 + 34.50}{2} \\ &= 34.53 \text{ cm}^3 \\ \text{b) } \text{mol } MnO_4^- &= 0.150 \times \frac{34.53}{1000} \\ &= 5.180 \times 10^{-3} \\ \text{c) } &\text{1 : 5 ratio } MnO_4^- : Fe^{2+} \\ &\text{so mol Fe} = 5 \times 5.180 \times 10^{-3} \\ &= 0.02590 \\ \text{d) } \text{moles of Fe} &= 0.02590 \\ \text{e) } \text{mass Fe} &= 0.02590 \times 55.85 \\ &= 1.445 \text{ g} \end{aligned}$$

**COPYRIGHT
PROTECTED**



Questions




Prepare your solutions

1. 50.0 cm³ of an iron(II) solution is titrated with 0.0200 mol dm⁻³ KMnO₄ solution. 18.50 cm³. The reaction is:



- Calculate the number of moles of MnO₄⁻ used in the titration.
- Calculate the number of moles of Fe²⁺ which reacted with MnO₄⁻.
- Find the concentration of Fe²⁺ ions in the 50.0 cm³ solution.

2. Iodine (I₂) and thiosulfate (S₂O₃²⁻) react together in the following reaction:



50.0 cm³ samples of an iodine solution were titrated with 0.120 mol dm⁻³ thiosulfate. The mean titre was 26.78 cm³.


- Calculate the number of moles of thiosulfate used in the mean titre.
- Calculate the number of moles of iodine required to react with this amount of thiosulfate.
- Calculate the concentration of iodine in the original solution.



High concentration

3. 3.25 g of a soluble iron(II) salt with the formula FeX₂ is dissolved in 1.00 dm³ water. The solution is then titrated with 0.0200 mol dm⁻³ KMnO₄, giving a mean titre of 25.65 cm³.

The following reaction occurs during the titration:



The identity of the anion represented by X in the formula FeX₂ can be determined from the formula of the salt. Determine the identity of the iron(II) salt.

- Calculate the number of moles of MnO₄⁻ used in the titration.
- Calculate the number of moles of Fe²⁺ which reacted with MnO₄⁻.
- Calculate the number of moles of Fe²⁺ ions in the 1.00 dm³ solution.
- Determine the relative formula mass of the iron salt, and so determine the identity of the salt.

4. A sample of impure iron with a mass of 6.00 g is reacted with excess nitric acid to produce iron(II) nitrate which is made up to 500 cm³. 25.0 cm³ samples of the solution are then titrated with 0.0200 mol dm⁻³ potassium permanganate, giving a mean titre of 23.93 cm³.

The following reaction occurs during the titration:



- Calculate the number of moles of MnO₄⁻ used in the titration.
- Calculate the number of moles of Fe²⁺ which reacted with MnO₄⁻.
- Calculate the number of moles of Fe²⁺ in the original sample.
- Calculate the mass of iron in the sample and hence determine the percentage of iron in the sample.

COPYRIGHT
PROTECTED





Drop the base

5. A 500 cm³ solution of hydrogen peroxide, H₂O₂, with a concentration of approximately 0.1 mol dm⁻³ is analysed using titration methods.

Step 1


8.40 g of potassium iodide is added to the hydrogen peroxide solution and the mixture is allowed to react.



Step 2


50.0 cm³ samples of the resulting solution are titrated with a solution of 0.150 mol dm⁻³ sodium thiosulfate, giving a mean titre of 26.70 cm³.

The following reaction occurs during the titration:



- Demonstrate that iodide ions are in excess of hydrogen peroxide in Step 1.
 - From the mean titre, deduce the number of moles of H₂O₂ in a 50.0 cm³ sample.
 - Calculate the concentration of H₂O₂ in the original 500 cm³ solution.
6. 6.03 g of a pure soluble iron(II) salt with the formula FeSO₄·nH₂O was dissolved in 250 cm³ of water. 25.0 cm³ samples were taken and titrated with 0.0150 mol dm⁻³ MnO₄⁻ solution, giving a mean titre of 12.50 cm³.

The following reaction occurs during the titration:



- Calculate the moles of MnO₄⁻ used in the titration.
- Calculate the moles of Fe²⁺ titrated.
- Calculate the moles of Fe²⁺ in the dissolved sample.
- Calculate the molar formula mass of the salt FeSO₄·nH₂O and thus determine n.

COPYRIGHT
PROTECTED



Unstructured redox titration

Unstructured calculations

Unstructured redox titration questions can be worth around 4–7 marks. They may be of a type which you haven't seen before.

When performing unstructured redox titration calculations, it can sometimes be helpful to have more information you have been given to the answer. Each step of the process will involve a calculation and putting them all together can be difficult with a lot of information.

Here is some advice for attempting unstructured redox titrations:


- Identify what you are trying to calculate – this could be:
 - relative simple – concentration, moles or mass
 - harder – purity, percentage by mass or relative formula mass
 - the most difficult – finding the charge on an ion or finding the formula of a compound
- You might find it helpful to write down all of the values you are given from the question to work out how all the pieces of information can be used together to find the answer.
- Convert values into the correct units so that you can use them in the equation.
- Make sure that you haven't missed a key step! In complex calculations it's much easier to miss a step than to have too many.

Examples

Example 1

A solution containing iodine is tested using redox titration methods.

50.0 cm³ samples of an iodine solution require 12.37 cm³ of 0.00700 mol dm⁻³ thiosulfate (S₂O₃²⁻) solution to reach the end point.




Find the concentration of iodine in the solution.

$$\begin{aligned} n(S_2O_3^{2-}) &= 0.00700 \times \frac{12.37}{1000} \\ &= 8.659 \times 10^{-5} \text{ mol} \\ n(I_2) &= \frac{1}{2} \times 8.659 \times 10^{-5} \\ &= 4.3295 \times 10^{-5} \text{ mol} \\ \text{conc } I_2 &= \frac{4.3295 \times 10^{-5}}{0.0500} \\ &= 8.659 \times 10^{-4} \text{ mol dm}^{-3} \end{aligned}$$

Example 2

A 1.00 g iron sample is reacted with dilute sulfuric acid until all of the iron dissolves, to form a 250 cm³ solution of iron(II) sulfate solution, FeSO₄.

25.0 cm³ samples of the solution are tested by redox titration methods with potassium permanganate, KMnO₄, requiring 31.03 cm³ of 0.00930 mol dm⁻³ permanganate solution to reach the end point.



Find the purity of the iron sample.

$$\begin{aligned} n(MnO_4^-) &= 0.00930 \times \frac{31.03}{1000} \\ &= 2.8858 \times 10^{-4} \text{ mol} \\ 5 : 1 \text{ ratio } Fe^{2+} : MnO_4^- \\ n(Fe^{2+}) &= 5 \times 2.8858 \times 10^{-4} \\ &= 0.001443 \text{ mol} \\ \text{conc } Fe^{2+} &= \frac{0.001443}{0.0250} \\ &= 0.05772 \text{ mol dm}^{-3} \\ \text{mass Fe} &= 0.05772 \times 0.0250 \times 55.85 \\ &= 0.805 \text{ g} \\ \text{purity Fe} &= \frac{0.805}{1.00} \times 100 \\ &= 80.5 \% \end{aligned}$$

INSPECTION COPY

COPYRIGHT
PROTECTED




Questions



Prepare your solutions

1. A solution containing iodine is tested using redox titration methods. 50.0 cm³ require 24.33 cm³ of 0.100 mol dm⁻³ thiosulfate (S₂O₃²⁻) solution to reach the end point.


The following reaction occurs during the titration:



Find the concentration of the iodine solution.

2. An iron(II) salt is reacted with dilute sulfuric acid until all of the iron dissolves, forming an iron(II) sulfate solution, FeSO₄. The solution is made up to 500 cm³. 20 cm³ sample is titrated by redox titration methods with potassium permanganate, KMnO₄, requiring 12.5 cm³ of 0.0200 mol dm⁻³ permanganate solution to reach the end point.

The following reaction occurs during the titration:




Find the mass of iron present in the sample.



High concentration

3. A soluble iron(II) salt has the formula FeX₂, where X represents one of Cl⁻, Br⁻ or I⁻. 10.0 g of the salt is dissolved in 100 cm³ water, and 20.0 cm³ samples are titrated against 0.0500 mol dm⁻³ potassium permanganate solution, giving a mean titre of 14.58 cm³.


The following reaction occurs during the titration:



Determine the identity of the X ion.

4. 50.0 cm³ of a solution containing Mn³⁺ reacts with a solid salt containing I⁻ ions, forming a 50.0 cm³ solution of product.

The iodine solution can then be titrated against a solution of thiosulfate, S₂O₃²⁻, during the titration:




The solution of thiosulfate is made by dissolving 5.00 g of Na₂S₂O₃•5H₂O in 500 cm³ of water. The thiosulfate is 34.55 cm³. Na₂S₂O₃•5H₂O has a relative formula mass of 248.0. Calculate the concentration of the original Mn³⁺ solution. (You will need to work out the equation for the first reaction.)



Drop the base

5. A sample of Fe(NO₃)₃ contains soluble, unreactive impurities. 5.64 g of the sample is dissolved in water, forming a solution containing Fe³⁺ ions. An excess of tin(II) chloride is added, reducing the Fe³⁺ ions to Fe²⁺ ions. The excess Sn²⁺ ions are removed by adding a solution of sodium hydroxide. The solution is made up to 500 cm³ and 25.0 cm³ of the solution is titrated against 0.00400 mol dm⁻³ potassium dichromate, K₂Cr₂O₇, giving a mean titre of 12.5 cm³.

Cr₂O₇²⁻ is reduced back to Fe³⁺ via the following half-equation:




Write the balanced equation for the reaction which occurs during the titration of iron nitrate in the sample.

COPYRIGHT
PROTECTED




6. Cuprite is an ore of copper. 832 mg of cuprite is added to concentrated hydrochloric acid forming a solution containing Cu^{2+} ions.

An excess of potassium iodide is added and the following reaction occurs:



The resulting solution is diluted to 500 cm^3 , and 25.0 cm^3 samples are titrated with sodium thiosulfate, giving a mean titre of 14.55 cm^3 .

During the titration, the following reaction occurs:



Calculate the percentage by mass of copper in the ore.



**COPYRIGHT
PROTECTED**



The most difficult redox titration

In past papers some of the most difficult redox titration questions have been worth 7-mark questions not uncommon. These questions outline a multistep titration procedure such as the percentage purity, the identity of an ion, or a balanced equation.

These questions are difficult because they require a lot of steps between the initial question and the final answer. The key is to recognise that most of the steps are exactly the same as more structured questions. You can still get most of the marks for showing the same set of steps involving moles, mass, and concentration, even if you've done many times. To get the final marks, you will need to pay very close attention to detail.

Top tips:

1. Start by working out the equations for any reactions taking place. This will often give you a balanced redox equation.
2. Identify all the quantities you are given in the equation, and what they relate to. A short list of what you know can be helpful.
3. Keep an eye out for dilution steps where 'samples' are taken or water is added. You would then work out the number of moles or the concentration of the original solution.


Examples

Example 1


A dicarboxylic acid with the formula $(\text{COOH})_2(\text{CH}_2)_n$ is analysed using titration methods with potassium permanganate, KMnO_4 . The formula $(\text{COOH})_2(\text{CH}_2)_n$ can be represented as H_2X in equations.

1.38 g of the acid is dissolved in water and made up to 900 cm^3 . 25.0 cm^3 samples gave a titre of 26.33 cm^3 when titrated against 0.0201 mol dm^{-3} KMnO_4 .

The half-equation for the oxidation of the acid is:









The half-equation for the reduction of permanganate is:



Find the identity of n.

Working out the redox equation:



$$\text{mol MnO}_4^- = 0.0201 \times \frac{26.33}{1000} = 5.2923 \times 10^{-3} \text{ mol}$$

$$\text{mol H}_2\text{X} = \frac{5 \times 5.2923 \times 10^{-3}}{2} = 0.01323 \text{ mol}$$

$$\text{r.f.m. H}_2\text{X} = \frac{1.38}{0.01323} = 104 \quad \leftarrow \text{r.f.m.} = \frac{\text{mass}}{\text{mol}}$$

$$\text{r.f.m. (CH}_2)_n = 104 - \underbrace{2 \times (12 + 16 \times 2 + 1)}_{\text{r.f.m. due to (COOH)}_2} = 14$$

$$n = 14 \div (12 + 2 \times 1) = 1$$

INSPECTION COPY


COPYRIGHT
PROTECTED



Example 2

0.510 g of sodium oxalate, $\text{Na}_2\text{C}_2\text{O}_4$, was titrated against a solution of 0.100 mol dm^{-3} potassium permanganate, KMnO_4 .

The half-equation for the permanganate ions is:



15.20 cm^3 of the potassium permanganate solution was required to reach the end point.

Calculate the ratio of MnO_4^- to $\text{C}_2\text{O}_4^{2-}$ in the overall equation, and hence work out the oxidation state of carbon in the carbon-containing product formed. (You may assume that both carbons end up with the same oxidation state.)

$$\begin{aligned} \text{mol MnO}_4^- &= 0.1 \times \frac{15.2}{1000} \\ &= 1.52 \times 10^{-3} \text{ mol} \end{aligned}$$

$$\text{r.f.m. Na}_2\text{C}_2\text{O}_4 = 134$$

$$\begin{aligned} \text{mol Na}_2\text{C}_2\text{O}_4 &= \frac{0.51}{134} \\ &= 3.8060 \times 10^{-3} \text{ mol} \end{aligned}$$

	$\text{C}_2\text{O}_4^{2-}$
mol ratio	3.8060×10^{-3}
whole number mol ratio	$\frac{3.8060 \times 10^{-3}}{1.52 \times 10^{-3}}$
	2.5
	5

$$\text{C}_2\text{O}_4^{2-} : \text{MnO}_4^- = 5 : 2$$

Mn changes oxidation state from +7 to

in MnO_4^- each O is -2, so Mn must be +7 to balance the -1 charge

Using the ratio 2 $\times$ MnO_4^- requires 10 e⁻. Mn is reduced by 5.

Therefore, each of the 5 $\text{C}_2\text{O}_4^{2-}$ ions in

Each carbon in $\text{C}_2\text{O}_4^{2-}$ must, therefore

Carbon's oxidation state in $\text{C}_2\text{O}_4^{2-}$ is +3. Its state at the end must be +4.

**COPYRIGHT
PROTECTED**




Questions



Prepare your solutions


- 5.68 g of an iron(II) salt is dissolved to form a 500 cm³ solution. 25.0 cm³ sample of 0.0250 mol dm⁻³ dichromate, Cr₂O₇²⁻, giving a mean titre of 18.90 cm³. The half-equations for the reaction are:



Use the half-equations to write a balanced equation for the redox reaction between dichromate ions (Cr₂O₇²⁻). Then, calculate the percentage by mass of iron in the salt.

- A 0.382 g sample of vanadium metal containing unreactive impurities is reacted with V²⁺ ions. 50.0 cm³ samples of the solution are analysed by redox titration with MnO₄⁻ solution, giving a mean titre of 21.1 cm³.

Write a balanced equation for the redox titration calculation, and find the percentage of vanadium in the metal.



High concentration

- A 500 cm³ solution with an unknown concentration of iron(II) sulfate is left in a beaker for several months, producing solution A. Iron(II) sulfate solution can oxidise over time.




Solution A is, therefore, a mixture of Fe(II) and Fe(III) ions.

The following method was used to find the percentage of iron ions which are Fe(II).

Step 1

25.0 cm³ samples of solution A are titrated with 0.0200 mol dm⁻³ KMnO₄ solution, giving a mean titre of 23.55 cm³.



Step 2

A 100 cm³ sample of solution A is passed over a reducing agent so that all of the Fe(III) becomes Fe(II). The solution is then made up to 200 cm³ with water.

Step 3

25.0 cm³ samples of the solution produced in step 2 are titrated with 0.0200 mol dm⁻³ KMnO₄ solution, giving a mean titre of 28.93 cm³.

Calculate the percentage of iron ions which are Fe²⁺ in solution A.

- An investigation is carried out to determine the reaction occurring when ferrate(VI) reacts with iodide ions.

Step 1

0.676 g of soluble salt barium ferrate(VI) is added to an alkaline solution, forming FeO₄²⁻ ions.

Step 2


An excess of acidified potassium iodide, KI, is added to the solution, which reacts with an iron species Feⁿ⁺, iodine (I₂) and Ba²⁺ ions. The resulting solution is made up to 250 cm³.

COPYRIGHT
PROTECTED



Step 3

20.0 cm³ samples of the resulting solution are titrated with 0.0100 mol dm⁻³ sodium tetrathionate, Na₂S₂O₃, giving a mean titre of 31.55 cm³. The equation for this redox reaction is



Determine the identity of the iron species Feⁿ⁺ and hence determine the balanced reaction between ferrate(VI) (FeO₄²⁻) and iodide ions, where I⁻ is converted to I₂.



Drop the base


5. A bottle of wine is tested for its alcohol concentration by performing the following steps.

Step 1

100 cm³ of wine is distilled to separate the ethanol, CH₃CH₂OH, from other components. The distillate is pipetted into a 500 cm³ volumetric flask which is made up to the line with distilled water.


Step 2

20.0 cm³ samples of the distillate are diluted to 100 cm³, and 25.0 cm³ of 0.119 mol dm⁻³ potassium dichromate(VI) solution is added and left to react to completion. Dichromate is in excess for titration.



Step 3

The samples are titrated with 0.109 mol dm⁻³ ammonium iron(II) sulfate solution as an indicator, to find the remaining concentration of Cr₂O₇²⁻.



A mean titre of 24.50 cm³ is recorded for the iron sulfate solution. Calculate the concentration of Cr₂O₇²⁻ in mol dm⁻³.

6. **Step 1**


2.00 g of anhydrous salt **A** containing molybdenum with the formula Na₂MoO₄·*x*H₂O is dissolved in distilled water.

Step 2

The solution is reacted with hydrochloric acid so that all of the molybdenum is in the form of Mo⁵⁺. The solution is made up to 200 cm³.

Step 3

50.0 cm³ samples of the solution of Mo⁵⁺ are then titrated against 0.103 mol dm⁻³ ceric(IV) sulfate solution, giving a mean titre of 20.05 cm³.



Step 4

A second sample containing 1.50 g of **A** is heated in a crucible, removing the water of crystallisation. The remaining anhydrous salt has a mass of 1.28 g.

Identify the formula of the salt, filling in the values of both *x* and *n*.


COPYRIGHT
PROTECTED



pH Titrations

1. A fizzy drink company is developing a new soft drink. The drink contains an organic acid.


The acid is monoprotic, which means it only ionises once:



A scientist tests the drink for the organic acid which is thought to have a concentration of $0.100 \text{ mol dm}^{-3}$.

The organic acid is titrated against NaOH, a strong base.

- a) Sketch a titration curve that you would expect on the axes when 40.0 cm^3 of $0.100 \text{ mol dm}^{-3}$ of the organic acid with a concentration of exactly $0.100 \text{ mol dm}^{-3}$.



- b) Results of the titration are shown below:

	Rough	1	2	
Start volume (cm^3)	0.00	0.10	0.05	0.00
End volume (cm^3)	33.35	31.15	31.15	33.35
Volume of NaOH (cm^3)				

- i) Complete the table and calculate the mean titre of NaOH to two decimal places.

.....

.....

.....

- ii) Find the actual concentration of the organic acid.

.....

.....

.....

INSPECTION COPY

COPYRIGHT
PROTECTED



- c) The organic acid has a pK_a of 3.13.
Find $[OH^-]$ for a $0.150 \text{ mol dm}^{-3}$ solution of the organic acid.
 $K_w = 1 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$.



- d) The drink manufacturer would like to ensure that the pH of the drink is kept constant. The scientist proposes making a 1.00 dm^3 buffer solution by adding 500 cm^3 of $0.100 \text{ mol dm}^{-3} \text{ NaOH}$.
- i) Explain how this buffer controls pH when a small amount of HCl is added. Include the equation for the buffer system equilibrium and explain how it works.




- ii) What concentration of the organic acid is required to produce a buffer of the volumes shown above?



**COPYRIGHT
PROTECTED**



2. Calcite is a mineral made from calcium carbonate, CaCO_3 .
A sample of calcite contains a small amount of quartz, made from silicon dioxide, SiO_2 .
The purity of calcium carbonate in calcite can be found by adding an excess of hydrochloric acid.
Calcium carbonate reacts with hydrochloric acid in the following reaction:



Silicon dioxide does not react with hydrochloric acid.
The amount of unreacted acid can then be determined by titration with sodium hydroxide.
Outline a method for an experiment to accurately calculate the purity of calcium carbonate in a sample of calcite.
You are given 100 cm³ of 0.150 mol dm⁻³ hydrochloric acid.



**COPYRIGHT
PROTECTED**



Redox Titrations


INSPECTION COPY

3. Rose gold is an alloy used to make jewellery. It contains the elements gold and copper. A rose gold ring with mass 0.945 g is added to sulfuric acid. The copper dissolves, but the gold does not dissolve.

The copper solution is added to an excess of potassium iodide and the following reaction occurs:



The solution is filtered to remove the solid copper iodide, and made up to 250 cm³ in a volumetric flask. Starch is added to 25.0 cm³ samples, which are titrated with 0.0100 mol dm⁻³ sodium thiosulfate. The reaction is:



The average titre is 37.20 cm³.

- a) Write the half-equation for copper in the first reaction.

.....

- b) Suggest why the solid copper iodide is removed by filtration before the titration.

.....

.....

.....

- c) State the role of starch in this process.

.....

.....

- d) Calculate the percentage by mass of copper in the rose gold ring.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

COPYRIGHT
PROTECTED



INSPECTION COPY



**COPYRIGHT
PROTECTED**



4. A pure salt with the formula M_2SO_3 contains ions of an unknown metal, M^+ . The salt is tested by redox titration using the following method:
- 1.24 g of the salt is dissolved in water.
 - The solution is made up to 200 cm^3 .
 - 25.0 cm^3 samples are titrated against an acidified solution of $0.0150\text{ mol dm}^{-3}$ $K_2Cr_2O_7$.

The end point was reached at 32.8 cm³.

The half-equations for the reaction are:



Construct the half-reaction for the reaction occurring in the titration between SO_3^{2-} metal ion and MnO_4^- is Na^+ .

Appendix: GCSE knowledge re

At GCSE, you will have learnt some very important things about acids, alkalis, titration and an essential foundation for the A Level material you need to know.

If you studied GCSE Chemistry, you will probably already be familiar with some aspects and will probably know your acids from your alkalis, and how they react together, the results tables used in titrations, and you may have used moles to calculate concentration in an experiment.

The next few pages are a quick summary of the GCSE material to jog your memory before moving on to the more difficult A Level material. If aren't feeling confident, spend some time building a strong foundation of knowledge and understanding before moving on to A Level material.

Acids and alkalis

Examples


Acids and alkalis can be dissolved in water to form solutions. Solutions of acids contain H^+ ions and alkalis contain OH^- ions.

Here are some examples of acids and alkalis, and the ions they form in solution:

Acids			Alkalis
Name	Formula	Ions in solution	Name
Hydrochloric acid	HCl	H^+ and Cl^-	Sodium hydroxide
Sulfuric acid	H_2SO_4	2H^+ and SO_4^{2-}	Potassium hydroxide
Phosphoric acid	H_3PO_4	3H^+ and PO_4^{3-}	Calcium hydroxide
Ethanoic acid	CH_3COOH	H^+ and CH_3COO^-	Magnesium hydroxide

Neutralisation

When an acid and an alkali are mixed together, a neutralisation reaction occurs, for example, when H^+ and OH^- react together:



This reaction is the general reaction for neutralisation reactions, because it occurs in all neutralisation reactions. The salt is formed from the remaining ions in the solution. For example, when HCl and NaOH are mixed, the salt formed is NaCl .

pH

The concentration of H^+ or OH^- ions in a solution determines its pH. pH is a scale, from 1 to 14, where lower numbers are more acidic. Here is a pH scale:



Indicator

You've probably used universal indicator at school to determine the pH of a solution. It is orange in acids, green at neutral and blue or purple in alkalis. There are many kinds of indicators, but they all have one similarity: they change colour depending on the pH.


INSPECTION COPY

COPYRIGHT
PROTECTED




Concentration of acid

Dissolving more acid in a solution will increase the number of H^+ ions in the solution.
 high concentration low concentration



A **dilute** acid can be made from a **concentrated** acid by adding more water.



Acidity

Strong acids ionise more than weak acids. Ionisation is when the ions in the acid 'split'. For example, when the strong acid HCl dissolves, almost all of the H^+ ions separate.

However, when the weak acid CH_3COOH dissolves, most of the H^+ ions remain bonded. So many free H^+ are in the solution. If the same amount of each acid is dissolved, the concentration of H^+ , and, therefore, a lower pH.

Strong acid				Weak acid	
For 100 HCl particles dissolved in solution the result is:				For 100 CH_3COOH dissolved in solution the result is:	
	HCl	$\rightarrow$	H^+	+	Cl^-
No. of particles	0		100		100
All of the acid is ionised, and there are many H^+ ions .				Only some are ionised and there are few H^+ ions .	

**COPYRIGHT
PROTECTED**



Concentrations and mole calculations in reactions

Concentration

The concentration of a solution depends on the amount of the solute and the volume

concentration represents either: the amount (in moles) dissolved in a volume (in dm^3) or the mass in a volume (in dm^3)

which can be calculated using:

$$\text{concentration} = \frac{\text{moles}}{\text{volume}}$$


units: mol dm^{-3}



Example

Dissolving 2 moles of sugar in 4 dm^3 of water:

$$\text{concentration} = \frac{\text{moles}}{\text{volume}} = \frac{2}{4} = 0.5 \text{ mol dm}^{-3}$$



Converting between mol dm^{-3} and g dm^{-3}

You can convert between mol dm^{-3} and g dm^{-3} just as you can convert between mol and g.

You need to use the relative molecular mass of the substance (known as M_r).

$$\text{mol} = \frac{\text{mass}}{M_r} \quad \text{or} \quad \text{mol dm}^{-3} = \frac{\text{g dm}^{-3}}{M_r}$$

Example

8 g of sodium hydroxide (NaOH) is dissolved in 2000 cm^3 of water.

NaOH has a M_r of 40

The concentration in g dm^{-3} is:

$$\text{concentration} = \frac{\text{grams}}{\text{volume}} = \frac{8}{2} = 4 \text{ g dm}^{-3}$$

The concentration in mol dm^{-3} is:

$$\frac{4 \text{ g dm}^{-3}}{40} = 0.1 \text{ mol dm}^{-3}$$

INSPECTION COPY


COPYRIGHT
PROTECTED



Titration method

Titration can be used to find the concentration of a solution by performing a reaction.

- An exact volume of the solution with unknown concentration is measured, and
- The solution with known concentration is added to the burette.
- The tap of the burette is opened carefully, and the known solution is added to the conical flask until the indicator changes colour, signifying the end point.
- The volume change of the known solution is recorded.
- The experiment is repeated.




Equipment	Purpose
Pipette (and filler)	Measuring an accurate volume of the unknown solution to go into the conical flask
Conical flask	Holds the solution with unknown concentration and the indicator
Indicator	Changes colour and shows that the reaction is complete
Funnel	Used to minimise spillage of the solution of known concentration
Burette	A very accurate and precise way of adding the solution of known concentration
White tile	A blank background to make it easier to see the colour change

**COPYRIGHT
PROTECTED**



Titration calculations

In the following reaction, an acid (HCl) reacts with an alkali (NaOH) to form a salt (NaCl) and water (H₂O). You should recognise this as a neutralisation reaction.



The premise of a titration experiment is that if you know the number of moles of acid, you can find the number of moles of base.

You will need to use the following equation:

$$\text{concentration} = \frac{\text{moles}}{\text{volume}}$$




Worked example

A solution of NaOH has an unknown concentration. HCl is added to 0.050 dm³ of the NaOH solution until the indicator changes colour to show that the solution is neutral.

0.020 dm³ of HCl with a concentration of 1.0 mol dm⁻³ was required to neutralise the NaOH.

This is the reaction:



Step 1: find the number of moles of HCl used

$$\begin{aligned} \text{moles HCl} &= \text{concentration} \times \text{volume} \\ &= 1.0 \times 0.020 \\ &= 0.020 \text{ mol} \end{aligned}$$

Step 2: find the number of moles of NaOH

$$\text{ratio of NaOH : HCl} = 1 : 1$$

$$\text{moles NaOH} = 0.020$$

Step 3: find the concentration of NaOH

$$\begin{aligned} \text{conc NaOH} &= \frac{\text{moles of NaOH}}{\text{volume}} \\ &= \frac{0.020}{0.050} \\ &= 0.40 \text{ mol dm}^{-3} \end{aligned}$$


Reduction and oxidation

Oxidation and reduction reactions

At GCSE you will have learnt that in reactions like the following, where substances are being oxidised:





... and that in the opposite of these reactions, where an element loses oxygen:



There is also a broader definition for redox reactions, in terms of elements gaining or losing electrons.

If an element **loses** electrons, then it is being oxidised.

If an element **gains** electrons, it is being reduced.




COPYRIGHT
PROTECTED



Quick quiz

1. Which ion is found in all acidic solutions?
2. What is the name of the type of reaction when an acid reacts with a base?
3. Complete all of the gaps in the following word and symbol equation:



4. On the pH scale, what does 7 represent?
5. Name two methods of measuring pH in a classroom.
6. 3 mol of an acid are dissolved in 9 dm³ of water.
Calculate the concentration of this solution in mol dm⁻³.
7. Name three pieces of practical equipment you might use during a titration.
8. A science teacher makes two solutions of acids:


Solution 1 contains 1 mol of sulfuric acid, a **strong acid**.

Solution 2 contains 1 mol of propanoic acid, a **weak acid**.

Both solutions are made using 500 cm³ of water.

Explain why **solution 1** has a higher concentration of hydrogen ions than **solution 2**.

9. a) Write the **balanced** symbol equation for the reaction between H₂SO₄ and NaOH.
b) How many molecules of water are formed when an excess of H₂SO₄ reacts with 10 molecules of NaOH?
10. Look at the following reaction:



Identify the elements which are:

- a) reduced
- b) oxidised
- c) neither reduced nor oxidised

COPYRIGHT
PROTECTED



Answers

Titration basics: results and uncertainties

1.

	Rough	1	2	3
Start volume (cm ³)	0.00	12.65	24.20	0.00
End volume (cm ³)	12.65	24.20	35.70	11.55
Volume of acid (cm ³)	12.65	11.55	11.50	11.55

2. a) 20.15 cm³
 b) 45.15 cm³
 c) 0.2 cm³
 d) 26.80 cm³

3.

	Rough	1	2	3
Start volume (cm ³)	0.80	0.20	0.55	0.50
End volume (cm ³)	17.10	17.35	17.30	17.35
Volume of acid (cm ³)	16.30	17.15	16.75	16.85

$$\frac{16.75 + 16.85}{2} = 16.80 \quad [\text{to 2 d.p.}]$$

4.

	Rough	1	2	3
Start volume (cm ³)	0.05	19.50	0.10	19.05
End volume (cm ³)	19.50	38.40	19.20	38.05
Volume of acid (cm ³)	19.45	18.90	19.10	19.00

$$\text{mean result} = \frac{18.90 + 18.95 + 18.90}{3} = 18.92 \text{ cm}^3 \quad [\text{to 2 d.p.}]$$

5.

	Rough	1	2	3
Start volume (cm ³)	0.00	0.55	0.40	0.15
End volume (cm ³)	43.75	43.10	43.05	43.05
Volume of acid (cm ³)	43.75	42.55	42.65	42.90

$$\text{Mean titre} = \frac{42.55 + 42.65}{2} = 42.60$$

$$\text{Percentage uncertainty} = \frac{0.1}{42.60} \times 100 = 0.23 \%$$

6. Reduces the concentration of the acid / some of the acid is neutralised
 More [acid] is required to reach the end point
 Mean titre is higher

INSPECTION COPY

COPYRIGHT
PROTECTED



Concentrations and dilution

- $500 \div 1000 = 0.5 \text{ dm}^3$ $\frac{3}{0.5} = 6.00 \text{ g dm}^{-3}$
- $200 \div 1000 = 0.2 \text{ dm}^3$ $0.2 \times 0.2 = 0.0400 \text{ moles}$
- $\text{vol} = \frac{\text{mol}}{\text{conc}} = \frac{2.5}{0.5} = 5.00 \text{ dm}^3$
- $\text{conc} = 2 \times \frac{25}{1000} = 0.100 \text{ mol dm}^{-3}$
- $10 \times \frac{0.4}{0.0125} = 320 \text{ cm}^3$
- $\text{conc} = 0.24 \times \frac{100}{2.5} = 9.6 \text{ mol dm}^{-3}$
- mol KI = 0.0205 mol
 $\text{conc KI} = 0.0205 \div \frac{200}{1000} = 0.102 \text{ mol dm}^{-3}$
- Concentration before NaCl added = $1 \times \frac{250}{1000} = 0.250 \text{ g dm}^{-3}$
 Concentration after NaCl added = $0.25 + 0.5 = 0.750 \text{ g dm}^{-3}$

Structured acid-base titration calculations

1.

	Rough	1	2	3
Start volume (cm ³)	0.00	16.20	0.10	16.15
End volume (cm ³)	16.20	32.25	16.15	32.25
Volume of acid (cm ³)	16.20	16.05	16.05	16.10

a) $\frac{16.05 + 16.05 + 16.10}{3} = 16.07 \text{ cm}^3$

b) $0.2 \times \frac{16.07}{1000} = 3 \times 10^{-3} \text{ mol}$

2.

	Rough	1	2	3
Start (cm ³)	0.00	0.00	23.85	0.00
End volume (cm ³)	24.40	23.85	47.65	23.80
Volume of acid (cm ³)	24.40	23.85	23.80	23.80

a) $\frac{23.85 + 23.80 + 23.80}{3} = 23.82 \text{ cm}^3$

b) $0.1 \times \frac{23.82}{1000} = 2.382 \times 10^{-3} \text{ mol}$

c) $2.382 \times 10^{-3} \times 2 = 4.76 \times 10^{-3} \text{ mol}$

- a) mol HCl = $\frac{13.7}{1000} \times 0.2 = 0.00274 \text{ mol dm}^{-3}$

b) mol KOH = 0.00274 mol [1 : 1 reaction KOH : HCl]

c) conc KOH = $0.00274 \div \frac{30}{1000} = 0.0913 \text{ mol dm}^{-3}$

- a) conc H₂SO₄ = $2.00 \times \frac{250}{1000} = 0.500 \text{ mol dm}^{-3}$

b) mol H₂SO₄ = $0.0192 \text{ mol dm}^{-3}$

c) mol KOH = $0.0192 \times 2 = 0.0384 \text{ mol}$ [2 : 1 reaction KOH : H₂SO₄]

d) conc KOH = $0.0384 \div \frac{80}{1000} = 0.480 \text{ mol dm}^{-3}$

**COPYRIGHT
PROTECTED**



5. a) Results table:

	Rough	1	2	3
Start volume (cm ³)	0.05	0.40	0.40	0.20
End volume (cm ³)	41.95	40.55	40.50	40.50
Volume of acid (cm ³)	41.90	40.15	40.10	40.30

$$\text{Mean titre} = \frac{40.15 + 40.10}{2}$$

$$= 40.13 \text{ cm}^3$$

b) mol HCl = $\frac{40.13}{1000} \times 0.206$

$$= 0.008268 \text{ mol dm}^{-3}$$

mol Ca(OH)₂ = $0.008268 \div 2$

$$= 0.0041334 \text{ mol} \quad [1 : 2 \text{ reaction Ca(OH)}_2 : \text{HCl}]$$

mol Ca(OH)₂ = $0.0041334 \div \frac{250}{1000}$

$$= 0.01653 \text{ mol dm}^{-3} \text{ [in 250 cm}^3 \text{ samples]}$$

c) conc Ca(OH)₂ = 0.01653 mol dm⁻³ [in 1 dm³ solution]

conc Ca(OH)₂ = $0.01653 \times \frac{1000}{200}$ [1000/200 is the dilution of 200 cm³]

$$= 0.0827 \text{ mol dm}^{-3} \quad [\text{In } 200 \text{ cm}^3 \text{ sample}]$$

d) M_r Ca(OH)₂ = 74.1

conc Ca(OH)₂ = 74.1×0.0827

$$= 6.13 \text{ g dm}^{-3}$$

Yes (it is above the required concentration)

6. a)

	Rough	1	2	3
Start volume (cm ³)	0.55	0.50	0.85	0.40
End volume (cm ³)	18.15	17.65	17.70	17.65
Volume of acid (cm ³)	18.65	17.15	17.85	17.25

mean titre = $\frac{17.15 + 17.25}{2}$

$$= 17.20 \text{ cm}^3$$

mol HCl = $\frac{17.20}{1000} \times 0.245$

$$= 0.004214 \text{ mol}$$

b) mol MOH = 0.004214 [1 : 1 ratio HCl : MOH]

mass MOH = $\frac{236.4}{1000}$

$$= 0.2364 \text{ g in } 200 \text{ cm}^3$$

$$= 0.02364 \text{ g in } 20 \text{ cm}^3$$

M_r MOH = $\frac{0.236}{0.004214}$

$$= 56.1 \quad [3 \text{ s.f.}]$$

COPYRIGHT
PROTECTED



Unstructured acid–base titration calculations

1. Results table:

	Rough	1	2	3
Start volume (cm ³)	0.00	18.70	0.65	17.65
End volume (cm ³)	18.70	35.45	17.65	34.70
Volume of acid (cm ³)	18.70	16.75	17.00	17.05

$$\text{mean titre} = \frac{17.00 + 17.05}{2}$$

$$= 17.03 \text{ cm}^3$$

$$\text{moles of HCl} = 0.150 \times \frac{17.03}{1000}$$

$$= 0.002555 \text{ mol}$$

1 : 1 ratio KOH : HCl

$$\text{moles of KOH} = 0.002555 \text{ mol}$$

$$\text{conc KOH} = 0.002555 \div \frac{50}{1000}$$

$$= 0.0511 \text{ mol dm}^{-3}$$

2. Results table:

	Rough	1	2	3
Start volume (cm ³)	0.45	0.55	0.00	0.40
End volume (cm ³)	32.55	31.50	31.55	31.35
Volume of acid (cm ³)	33.00	30.95	31.55	30.95

$$\text{mean titre} = \frac{30.95 + 30.95}{2}$$

$$= 30.95 \text{ cm}^3$$

$$\text{moles KOH} = 0.130 \times \frac{25.0}{1000}$$

$$= 0.00325 \text{ mol}$$

2 : 1 ratio H₂SO₄ : KOH

$$\text{moles H}_2\text{SO}_4 = \frac{0.00325}{2}$$

$$= 0.001625$$

$$\text{conc H}_2\text{SO}_4 = 0.001625 \div \frac{30.95}{1000}$$

$$= 0.0525 \text{ mol dm}^{-3}$$

$$\text{3. mol H}_2\text{SO}_4 = 0.100 \times \frac{30.58}{1000}$$

$$= 0.003058 \text{ mol}$$

$$\text{mol MOH in 25 cm}^3 = 0.003058 \times 2$$

$$= 0.006116 \text{ mol}$$

$$\text{mol MOH in 250 cm}^3 = 0.006116 \times \frac{100}{25}$$

$$= 0.02446 \text{ mol}$$

$$M_r \text{ MOH} = \frac{\text{mass}}{\text{mol}}$$

$$= \frac{1.37}{0.02446}$$

$$= 55.9$$

MOH is KOH (M

INSPECTION COPY

COPYRIGHT
PROTECTED



4. Balanced equation:
 $2\text{HCl} + \text{K}_2\text{CO}_3 \rightarrow 2\text{KCl} + \text{CO}_2 + \text{H}_2\text{O}$

	Rough	1	2	3
Start volume (cm ³)	0.55	0.30	0.05	0.15
End volume (cm ³)	41.90	41.10	41.10	41.05
Volume of acid (cm ³)	42.45	40.80	41.05	40.90

mean titre = $\frac{40.80 + 40.90}{2}$

= 40.85 cm³

mol HCl = $0.125 \times \frac{40.85}{1000}$

= 0.005106 mol

mol K₂CO₃ = $\frac{0.005106}{2}$

= 0.002553 mol

mol K₂CO₃ in 500 cm³ = $0.002553 \times \frac{500}{25}$

= 0.05106

M_r K₂CO₃ = 138.2

mass K₂CO₃ = 138.2×0.05106

= 7.056 g

% purity = $\frac{7.056}{9.33} \times 100$

= 75.6 %

5. $\text{KOH} + \text{HNO}_3 \rightarrow \text{KNO}_3 + \text{H}_2\text{O}$

mol HNO₃ = $0.200 \times \frac{21.80}{1000}$

= 0.00436 mol

mol KOH in total = 0.00436 mol

mol KOH in A = $0.143 \times \frac{10}{1000}$

= 0.00143 mol

mol KOH = mol in total – mol in A

= 0.00436 – 0.00143

= 0.00293 mol

conc B = $0.00293 \div \frac{10}{1000}$

= 0.293 mol dm⁻³

6. mol H₂SO₄ = $0.150 \times \frac{28.85}{1000}$

= 0.0043275 mol

M_r Ba(OH)₂ = 171.7

mol Ba(OH)₂ in 250 cm³ = $2.86 \div 171.7$

= 0.016657 mol

mol Ba(OH)₂ in 25 cm³ = 0.0016657 mol

mol H₂SO₄ which reacts with NaOH = 0.0043275 – 0.0016657

= 0.0026618 mol

mol NaOH in 25 cm³ = 0.0026618 mol

= 0.0026618 mol

mol NaOH in 250 cm³ = 0.026618 mol

M_r NaOH = 40.0

mass NaOH = 40.0×0.053236

= 2.13 g

mass of impurity = $5.34 - (2.86 + 2.13) = 0.35 \text{ g}$

**COPYRIGHT
PROTECTED**



'Back' titrations

1. a) $\text{mol KOH} = 0.1 \times \frac{17.88}{1000} = 0.001788 \text{ mol}$
 $\text{mol HCl in } 50 \text{ cm}^3 = 0.001788 \text{ mol}$
 b) $\text{mol HCl in } 250 \text{ cm}^3 = 0.001788 \times \frac{250}{50} = 0.00894 \text{ mol}$
 (from titration)
 c) $\text{original mol HCl} = \frac{200}{1000} \times 0.1 = 0.02 \text{ mol}$
 (in stock solution)
 $\text{mol HCl reacted} = 0.02 - 0.00894 = 0.01106 \text{ mol}$
 d) $\text{mol CaCO}_3 = \frac{0.01106}{2} = 0.00553 \text{ mol}$
 $M_r \text{ CaCO}_3 = 100.1$
 $\text{mass CaCO}_3 = 100.1 \times 0.00553 = 0.554 \text{ g}$

2. a) $\text{mol KOH} = 0.100 \times \frac{32.45}{1000} = 0.003245 \text{ mol}$
 b) $\text{mol HCl in } 20.0 \text{ cm}^3 = 0.003245 \text{ mol}$
 $\text{mol HCl in } 100 \text{ cm}^3 = 0.003245 \times \frac{100}{20} = 0.016225 \text{ mol}$
 c) $\text{mol HCl reacted} = 0.0250 - 0.016225 = 0.008775 \text{ mol}$
 d) $\text{mol NH}_3 \text{ in } 30 \text{ cm}^3 = 0.008775 \text{ mol}$
 $\text{mol NH}_3 \text{ in } 3.00 \text{ dm}^3 = 0.008775 \times (3 \div \frac{30}{1000}) = 0.8775 \text{ mol}$

3. a) $3\text{M(OH)}_2 + 2\text{H}_3\text{PO}_4 \rightarrow \text{M}_3(\text{PO}_4)_2 + 6\text{H}_2\text{O}$
 b) Results table:

	rough	1	2	3
Sample volume (cm ³)	0.55	0.15	0.05	0.20
End volume (cm ³)	40.00	39.00	38.95	39.00
Volume of acid (cm ³)	39.45	38.85	38.90	38.80

- Mean titre $= \frac{38.85 + 38.90 + 38.80}{3} = 38.85 \text{ cm}^3$
- $\text{mol KOH} = \frac{38.85}{1000} \times 0.130 = 0.0050505 \text{ mol}$
- $\text{mol H}_3\text{PO}_4 \text{ in } 25 \text{ cm}^3 = \frac{0.0050505}{3} = 0.0016835 \text{ mol}$
- $\text{mol H}_3\text{PO}_4 \text{ in } 250 \text{ cm}^3 = 0.016835 \text{ mol}$
- $\text{original mol H}_3\text{PO}_4 = \frac{200}{1000} \times 0.15 = 0.0300$
- $\text{mol H}_3\text{PO}_4 \text{ reacted} = 0.03 - 0.016835 = 0.013165 \text{ mol}$
- $\text{mol M(OH)}_2 = 0.013165 \times 2 = 0.02633 \text{ mol}$
- $M_r \text{ M(OH)}_2 = 1.15 \div 0.0197475 = 58.2$
- c) Mg(OH)_2

**COPYRIGHT
PROTECTED**



4. This would be an 8-mark question in an exam. Marks are shown by ✓.

Part 1: Calculation of solid so that acid is in excess

Recognition that nitric acid must be in excess ✓

$$\text{mol HNO}_3 = 0.150 \times \frac{100}{1000} = 0.0150 \text{ mol} \quad \checkmark$$

Ratio from reaction equation is 2 : 1 $\text{HNO}_3 : \text{Na}_2\text{CO}_3$

so mol Na_2CO_3 is less than 0.0075 mol ($0.0150 \times \frac{1}{2}$) ✓

$$\text{Mass of Na}_2\text{CO}_3 \text{ weighed out} < 0.0075 \times (23 \times 2 + 12 + 3 \times 16) = < 1.74 \text{ g}$$

Part 2: Method

Use pipette to measure 100 cm³ of acid and add weighed solid to acid ✓

Titrate unreacted acid with 0.100 mol dm⁻³ KOH and record titre when indicator

Part 3: Calculation

$$\text{mol Na}_2\text{CO}_3 \cdot 7\text{H}_2\text{O} = 0.0150 - \left(\frac{\text{mean titre}}{2} \times 0.1 \right) \quad \checkmark$$

$$M_r = \frac{\text{mass of sodium carbonate heptahydrate}}{\text{moles of sodium carbonate heptahydrate}} \quad \checkmark$$

$$\begin{aligned} 5. \quad \text{mol NaOH} &= 0.1 \times \frac{20.08}{1000} \\ &= 0.002008 \text{ mol} \\ \text{mol HCl in } 50 \text{ cm}^3 &= 0.002008 \text{ mol} \\ \text{mol HCl in } 250 \text{ cm}^3 &= 0.002008 \times \frac{250}{50} \\ &= 0.01004 \text{ mol} \\ \text{original mol HCl} &= \frac{150}{1000} \times 0.24 \\ &= 0.0360 \text{ mol} \\ \text{mol HCl reacted} &= 0.0360 - 0.01004 \\ &= 0.02596 \text{ mol} \\ \text{mol MgCO}_3 \cdot x\text{H}_2\text{O} &= 0.02596 \div 2 \\ &= 0.01298 \text{ mol} \\ \text{mass MgCO}_3 \cdot x\text{H}_2\text{O} &= 144.14 - 142.34 \\ &= 1.80 \text{ g} \\ M_r \text{ MgCO}_3 \cdot x\text{H}_2\text{O} &= \frac{1.80}{0.01298} \\ &= 138.7 \\ M_r x\text{H}_2\text{O} &= 138.7 - 84.3 \\ &= 54.4 \\ x &= 54.4 \div 18 \\ &= 3 \end{aligned}$$

Formula is $\text{MgCO}_3 \cdot 3\text{H}_2\text{O}$

$$\begin{aligned} 6. \quad \text{mol NaOH} &= \frac{41.30}{1000} \times 0.100 \\ &= 0.00413 \text{ mol} \\ \text{mol H}_2\text{SO}_4 \text{ in } 50 \text{ cm}^3 &= 0.00413 \div 2 \\ &= 0.002065 \text{ mol} \\ \text{mol H}_2\text{SO}_4 \text{ in } 500 \text{ cm}^3 &= 0.02065 \text{ mol} \end{aligned}$$

$$\begin{aligned} M_r \text{ BaSO}_4 &= 233.4 \\ \text{mol BaSO}_4 &= \frac{3.05}{233.4} \end{aligned}$$

$$= 0.01307 \text{ mol}$$

$$\text{mol H}_2\text{SO}_4 \text{ reacted with Ba(OH)}_2 = 0.01307 \text{ mol}$$

$$\text{original mol H}_2\text{SO}_4 = \frac{250}{1000} \times 0.200$$

$$= 0.0500 \text{ mol}$$

$$\text{mol H}_2\text{SO}_4 \text{ reacted with NaOH} = \text{original mol} - \text{remaining mol} - \text{mol reacted}$$

$$= 0.0500 - 0.01307 - 0.02065$$

$$= 0.01628 \text{ mol}$$

$$\text{mol NaOH} = 0.01628 \times 2$$

$$= 0.03256 \text{ mol}$$

$$\text{conc NaOH} = 0.03256 \div \frac{200}{1000}$$


$$= 0.163 \text{ mol dm}^{-3}$$

**COPYRIGHT
PROTECTED**




Titration curves / indicators

- C
 - D
 - A
- bromothymol blue
 - bromocresol green
 - blue to colourless
- blue (phenolphthalein and thymolphthalein would both be colourless at this pH; b giving the solution its colour)
- Graph should be a strong acid–strong base curve with an equivalence point at 35



- Graph should be a strong acid–weak base curve with an equivalence point at 20



- bromothymol blue and phenolphthalein
 - methyl orange and bromothymol blue
 - bromothymol blue

COPYRIGHT
PROTECTED



Strong acids and bases

1. a) $\text{pH} = -\log_{10}(0.00100) = 3.00$
 b) $\text{conc} = 10^{-2.3} = 0.00501 \text{ mol dm}^{-3}$
 c) $\text{pH} = -\log_{10}(2 \times 0.000400) = 3.10$

2. a) $[\text{H}^+] = \frac{1.00 \times 10^{-14}}{0.00160}$
 $= 6.25 \times 10^{-13} \text{ mol dm}^{-3}$
 $\text{pH} = -\log_{10}(6.25 \times 10^{-13})$
 $= 11.2$

- b) $[\text{H}^+] = \frac{1.00 \times 10^{-14}}{0.000810}$
 $= 1.23 \times 10^{-12} \text{ mol dm}^{-3}$
 $\text{pH} = -\log_{10}(1.23 \times 10^{-12})$
 $= 11.9$

- c) $[\text{H}^+] = \frac{1.00 \times 10^{-14}}{2 \times 0.0000451}$
 $= 1.11 \times 10^{-10} \text{ mol dm}^{-3}$
 $\text{pH} = -\log_{10}(1.11 \times 10^{-10})$
 $= 9.96$

3. $[\text{H}^+] = 10^{-11.7}$
 $= 1.995 \times 10^{-12}$
 $[\text{H}^+] \times [\text{OH}^-] = 1.00 \times 10^{-14}$
 $[\text{OH}^-] = \frac{1.00 \times 10^{-14}}{[\text{H}^+]}$
 $= \frac{1.00 \times 10^{-14}}{1.995 \times 10^{-12}}$
 $= 5.01 \times 10^{-3} \text{ mol dm}^{-3}$

4. $[\text{H}^+] = 2 \times 0.000520$
 $= 1.04 \times 10^{-3} \text{ mol dm}^{-3}$
 $[\text{OH}^-] = \frac{1.00 \times 10^{-14}}{[\text{H}^+]}$
 $= \frac{1.00 \times 10^{-14}}{1.04 \times 10^{-3}}$
 $= 9.62 \times 10^{-12} \text{ mol dm}^{-3}$

5. $\text{mol HCl} = 0.01 \times \frac{500}{1000}$
 $= 5 \times 10^{-3} \text{ mol}$
 $\text{mol NaOH} = 0.03 \times \frac{150}{1000}$
 $= 4.5 \times 10^{-3} \text{ mol}$
 $\text{new mol HCl} = 5 \times 10^{-3} - 4.5 \times 10^{-3}$
 $= 5 \times 10^{-4} \text{ mol}$
 $\text{new conc HCl} = 5 \times 10^{-4} \div \frac{650}{1000}$
 $= 7.692 \times 10^{-4} \text{ mol dm}^{-3}$
 $\text{pH} = -\log_{10}(7.692 \times 10^{-4})$
 $= 3.11$

6. a) $[\text{H}^+] = 10^{-7}$
 $= 1.0 \times 10^{-7} \text{ mol dm}^{-3}$
 $[\text{OH}^-] = \frac{2.88 \times 10^{-14}}{1.0 \times 10^{-7}}$
 $= 2.88 \times 10^{-8} \text{ mol dm}^{-3}$
 $[\text{H}^+] > [\text{OH}^-]$, so **acidic**
 b) $[\text{OH}^-] = 0.0012$
 $= 1.2 \times 10^{-4} \text{ mol dm}^{-3}$
 $[\text{H}^+] = \frac{2.88 \times 10^{-14}}{2.1 \times 10^{-4}}$
 $= 1.371 \times 10^{-11} \text{ mol dm}^{-3}$
 $\text{pH} = -\log_{10}(1.371 \times 10^{-11})$
 $= 10.86$

COPYRIGHT
PROTECTED



Weak acids

- $-\log_{10}(1.42 \times 10^{-5}) = 4.85$
 - $-\log_{10}(4.98 \times 10^{-7}) = 6.30$
 - $10^{-1.03} = 9.33 \times 10^{-2}$
 - $10^{-3.12} = 7.59 \times 10^{-4}$

- $K_a = \frac{[H^+][SCN^-]}{[HSCN]}$
 - $[H^+] = [SCN^-]$
 $K_a = \frac{[H^+]^2}{[HSCN]}$
 $[H^+] = \sqrt{[HSCN] \times K_a}$
 $= \sqrt{0.0100 \times 1.1 \times 10^{-4}}$
 $= 1.05 \times 10^{-3} \text{ mol dm}^{-3}$
 $pH = -\log_{10}(0.00105)$
 $= 0.963$

- $K_a = 10^{-3.29}$
 $= 5.129 \times 10^{-4}$
 $= \frac{[H^+][NO_2^-]}{[HNO_2]}$
 $[NO_2^-] = [H^+]$
 $K_a = \frac{[NO_2^-]^2}{[HNO_2]}$
 $[NO_2^-] = \sqrt{[HNO_2] \times K_a}$
 $= \sqrt{0.0452 \times 5.129 \times 10^{-4}}$
 $= 4.81 \times 10^{-3} \text{ mol dm}^{-3}$

- $[H^+] = 10^{-5.79}$
 $= 1.622 \times 10^{-6} \text{ mol dm}^{-3}$
 $K_a = \frac{[H^+][C_6H_5O^-]}{[C_6H_5OH]}$
 $[H^+] = [C_6H_5O^-]$
 $K_a = \frac{[H^+]^2}{[C_6H_5OH]}$
 $K_a = \frac{(1.622 \times 10^{-6})^2}{1.0 \times 10^{-3}}$
 $= 2.63 \times 10^{-13}$
 $pK_a = 12.58$

- Ethanoic acid dissociates less in DMSO than in water.
 - $K_a(\text{DMSO}) = 10^{-12.3}$

$$K_a = \frac{[H^+][CH_3COO^-]}{[CH_3COOH]}$$

$$[H^+] = [CH_3COO^-]$$

$$K_a = \frac{[H^+]^2}{[CH_3COOH]}$$

$$[H^+] = \sqrt{[CH_3COOH] \times K_a}$$

$$= \sqrt{0.175 \times 5.0119 \times 10^{-13}}$$

$$= 2.962 \times 10^{-7} \text{ mol dm}^{-3}$$

$$K_a(\text{water}) = 10^{-4.76}$$

$$= 1.738 \times 10^{-5}$$

$$[CH_3COOH] = \frac{[H^+]^2}{K_a}$$

$$= \frac{(2.962 \times 10^{-7})^2}{1.738 \times 10^{-5}}$$

$$= 5.04 \times 10^{-9} \text{ mol dm}^{-3}$$

- $K_a = \frac{[H^+][H_2O]}{[H_3O^+]}$
 - $[H_2O] = \frac{1000}{18}$
 $= 55.56 \text{ mol dm}^{-3}$
 - $K_a = \frac{10^{-14}}{55.56}$
 $= 1.8 \times 10^{-16}$
 $pK_a = -\log(1.8 \times 10^{-16})$
 $= 15.7$

COPYRIGHT
PROTECTED



Buffers

- CHONa
 - NaCN⁻

- Equilibrium: $\text{CF}_3\text{COOH} \rightleftharpoons \text{CF}_3\text{COO}^- + \text{H}^+$
 Added HCl: CF_3COO^- reacts / uses up H^+ to form more CF_3COOH
 Equilibrium shifts left
 Added NaOH: H^+ reacts with added OH^- / more CF_3COO^- forms
 Equilibrium shifts right

- $$[\text{H}^+] = K_a \times \frac{[\text{HA}]}{[\text{A}^-]}$$

$$K_a = 10^{-8}$$

$$[\text{HA}] = 0.150 \times \frac{200}{1000} = 0.0300 \text{ mol dm}^{-3}$$

$$[\text{A}^-] = 0.0450 \times \frac{800}{1000} = 0.0360 \text{ mol dm}^{-3}$$

$$[\text{H}^+] = 9.120 \times 10^{-8} \times \frac{0.0300}{0.0360} = 7.6 \times 10^{-8}$$

$$\text{pH} = 7.12$$

- HF reacts with NH_3 to form F^- (or NH_4F)
 This means there is a mixture of leftover weak acid (because it was in excess) and its salt
 $\text{HF} \rightleftharpoons \text{F}^- + \text{H}^+$
 When:

acid added: F^- reacts with added H^+ / more HF forms

alkali added: H^+ reacts with added OH^- / more F^- forms

Equilibrium shifts left when acid is added, equilibrium shifts right when alkali is added

- $\text{CH}_3\text{CH}(\text{OH})\text{COOH}$ in excess
 - $[\text{H}^+] = 1 \times 10^{-4} \text{ mol dm}^{-3}$

$$K_a = 10^{-3.86} = 1.3804 \times 10^{-4}$$

Assume $[\text{KOH}] = [\text{CH}_3\text{CH}(\text{OH})\text{COO}^-]$

Assume $[\text{CH}_3\text{CH}(\text{OH})\text{COOH}] = [\text{CH}_3\text{CH}(\text{OH})\text{COOH}] - [\text{KOH}]$

$$[\text{H}^+] = K_a \times \frac{[\text{CH}_3\text{CH}(\text{OH})\text{COOH}] - [\text{KOH}]}{[\text{KOH}]}$$

$$[\text{CH}_3\text{CH}(\text{OH})\text{COOH}] - [\text{KOH}] = [\text{KOH}] \times \frac{[\text{H}^+]}{K_a}$$

$$[\text{CH}_3\text{CH}(\text{OH})\text{COOH}] = [\text{KOH}] \times \frac{[\text{H}^+]}{K_a} + [\text{KOH}]$$

$$= 0.05 \times \frac{1 \times 10^{-4}}{1.3804 \times 10^{-4}} + 0.05$$

$$= 0.08622 \text{ mol dm}^{-3}$$

$$\text{mol CH}_3\text{CH}(\text{OH})\text{COOH} = 0.08622 \times \frac{400}{1000} = 0.3449 \text{ mol}$$

$$M_r \text{ CH}_3\text{CH}(\text{OH})\text{COOH} = 77.0$$

$$\text{mass CH}_3\text{CH}(\text{OH})\text{COOH} = 26.6 \text{ g}$$

- Desired

$[\text{H}^+] =$

$=$

$K_a =$

$K_a =$

$=$

$[\text{HA}] =$

$=$

$[\text{A}^-] =$

$=$

$=$

current

current

addition

Conjugate

$M_r \text{ C}_5\text{H}_5\text{NO}_2$

mass C

INSPECTION COPY

COPYRIGHT
PROTECTED



Structured redox titrations

1. a) $\text{mol MnO}_4^- = 0.02 \times \frac{18.5}{1000} = 3.7 \times 10^{-4} \text{ mol}$
 b) 1 : 5 ratio
 $\text{mol Fe}^{2+} = 5 \times 3.7 \times 10^{-4} = 1.85 \times 10^{-3} \text{ mol}$
 c) $\text{conc Fe}^{2+} = 1.85 \times 10^{-3} \div \frac{50}{1000} = 0.037 \text{ mol dm}^{-3}$
2. a) $\text{mol S}_2\text{O}_3^{2-} = 0.12 \times \frac{10}{1000} = 1.2 \times 10^{-3} \text{ mol}$
 b) $\text{mol I}_2 = \frac{3.2136 \times 10^{-3}}{2} = 1.6068 \times 10^{-3} \text{ mol}$
 c) $\text{conc I}_2 = 1.6068 \times 10^{-3} \div \frac{50}{1000} = 0.0321 \text{ mol dm}^{-3}$
3. a) $\text{mol MnO}_4^- = 0.02 \times \frac{25.65}{1000} = 5.13 \times 10^{-4} \text{ mol}$
 b) 1 : 5 ratio
 $\text{mol Fe}^{2+} = 5.13 \times 10^{-4} \times 5 = 0.002565 \text{ mol}$
 c) $\text{mol Fe}^{2+} = 0.002565 \times \frac{1000}{100} = 0.02565 \text{ mol}$
 d) r.f.m. $\text{FeX}_2 = 3.25 \div 0.02565 = 126.7$
 $\text{r.f.m. X} = \frac{126.7 - 58.8}{2} = 35.5$
 X is Cl, so salt is FeCl_2
4. a) $\text{mol MnO}_4^- = 9.572 \times 10^{-4} \text{ mol}$
 b) 1 : 5 ratio
 $\text{mol Fe}^{2+} = 5 \times 9.572 \times 10^{-4} = 4.786 \times 10^{-3} \text{ mol}$
 c) $\text{mol Fe}^{2+} = 4.786 \times 10^{-3} \times \frac{500}{25} = 0.09572 \text{ mol}$
 d) $\text{mass Fe} = 0.09572 \times 55.8 = 5.34 \text{ g}$
 $\text{purity Fe} = \frac{5.34}{6} \times 100 = 89 \%$
5. a) $\text{mol KI} = \frac{8.4}{166} = 0.0506$
 $\text{mol H}_2\text{O}_2 = \sim 0.04$
 $\text{mol I}^- > 2 \times \text{mol H}_2\text{O}_2$
 $\text{mol Na}_2\text{S}_2\text{O}_3 = 0.150$
 $\text{mol I}_2 = \frac{4.017 \times 10^{-3}}{3} = 2.0085 \times 10^{-3} \text{ mol}$
 $\text{mol H}_2\text{O}_2 = 2.0085 \times 10^{-3} \text{ mol}$
 $\text{conc H}_2\text{O}_2 = 0.0200$
 $\text{conc H}_2\text{O}_2 = 0.0200$
 $\text{conc H}_2\text{O}_2 = 0.0402$
6. a) $\text{mol MnO}_4^- = 0.0150$
 $\text{mol Fe}^{2+} = 5 \times 4.32 \times 10^{-4} = 0.00216$
 $\text{mol Fe}^{2+} = 0.0216$
 $\text{r.f.m. FeSO}_4 \cdot n\text{H}_2\text{O} = 6$
 $n = 7$

COPYRIGHT
PROTECTED




Unstructured redox titrations

- $\text{mol S}_2\text{O}_3^{2-} = 0.1 \times \frac{24.33}{1000} = 2.433 \times 10^{-3} \text{ mol}$
 $2 : 1 \text{ ratio of S}_2\text{O}_3^{2-} : \text{I}_2$
 $\text{mol I}_2 = 2.433 \times 10^{-3} \div 2 = 1.2165 \times 10^{-3} \text{ mol}$
 $\text{conc I}_2 = 1.2165 \times 10^{-3} \div \frac{50}{1000} = 0.0243 \text{ mol dm}^{-3}$
- $\text{mol MnO}_4^- = 50 \times \frac{4.00}{1000} = 2.2275 \times 10^{-4} \text{ mol}$
 $5 : 1 \text{ ratio of Fe}^{2+} : \text{MnO}_4^-$
 $\text{mol Fe}^{2+} \text{ in } 25 \text{ cm}^3 = 5 \times 2.2275 \times 10^{-4} = 1.11375 \times 10^{-3} \text{ mol}$
 $\text{mol Fe}^{2+} \text{ in } 500 \text{ cm}^3 = 1.11375 \times 10^{-3} \times \frac{500}{20}$
 $= 0.02784 \text{ mol}$
 $\text{mass Fe in sample} = 0.02784 \times 55.8 = 1.55 \text{ g}$
- $\text{mol MnO}_4^- = 0.050 \times \frac{14.85}{1000} = 7.425 \times 10^{-4} \text{ mol}$
 $5 : 1 \text{ ratio of Fe}^{2+} : \text{MnO}_4^-$
 $\text{mol Fe}^{2+} \text{ in } 20 \text{ cm}^3 = 5 \times 7.425 \times 10^{-4} = 3.7125 \times 10^{-3} \text{ mol}$
 $\text{mol Fe}^{2+} \text{ in } 100 \text{ cm}^3 = 3.7125 \times 10^{-3} \times \frac{100}{20}$
 $= 0.01856 \text{ mol}$
 $\text{r.f.m. FeX}_2 = 2.74 \div 0.01856 = 147.8$
 $\text{r.f.m. X} = \frac{147.8 - 56}{2} = 45.9$
 X is NO_2
- $\text{mol Na}_2\text{S}_2\text{O}_3 \text{ in solution} = 5 \div 248 = 0.02016 \text{ mol}$
 $\text{conc of S}_2\text{O}_3^{2-} \text{ in solution} = 0.02016 \div \frac{500}{1000} = 0.04032 \text{ mol dm}^{-3}$
 $\text{mol S}_2\text{O}_3^{2-} \text{ in titre} = 0.04032 \times \frac{34.55}{1000} = 1.393 \times 10^{-3} \text{ mol}$
 $\text{mol I}_2 = \frac{1.393 \times 10^{-3}}{2} = 6.966 \times 10^{-4} \text{ mol}$
 $\text{Equation: } 2\text{Mn}^{3+} + 2\text{I}^- \rightarrow 2\text{Mn}^{2+} + \text{I}_2$
 $\text{mol Mn}^{3+} = 6.966 \times 10^{-4} \text{ mol} \times 2 = 1.3932 \times 10^{-3} \text{ mol}$
 $\text{conc I}_2 = 1.3932 \times 10^{-3} \div \frac{50}{1000} = 0.0279 \text{ mol dm}^{-3}$
- $\text{Cr}_2\text{O}_7^{2-} + 6\text{Fe}^{2+} + 14\text{H}^+ \rightarrow 2\text{Cr}^{3+} + 6\text{Fe}^{3+} + 7\text{H}_2\text{O}$
 $\text{mol Cr}_2\text{O}_7^{2-}$
 $\text{mol Fe}^{2+} \text{ (in } 25 \text{ cm}^3)$
 $\text{mol Fe}^{2+} \text{ (in } 500 \text{ cm}^3)$
 $\text{mol Fe(NO}_3)_3$
 $\text{r.f.m. Fe(NO}_3)_3$
 $\text{mass Fe(NO}_3)_3$
 $\text{purity Fe(NO}_3)_3$
- $\text{mol S}_2\text{O}_3^{2-}$
 $\text{mol I}_2 \text{ in } 25 \text{ cm}^3$
 $\text{mol I}_2 \text{ in } 500 \text{ cm}^3$
 mol Cu^{2+}
 mass Cu^{2+}
 dry mass




COPYRIGHT
PROTECTED



The most difficult redox titrations



mol $\text{Cr}_2\text{O}_7^{2-}$ = $0.0250 \times \frac{18.90}{1000}$
 = $4.725 \times 10^{-4} \text{ mol}$
 mol Fe^{2+} in 25 cm^3 = $6 \times 4.725 \times 10^{-4}$
 = $2.835 \times 10^{-3} \text{ mol}$
 mol Fe^{2+} in 500 cm^3 = $2.835 \times 10^{-3} \times \frac{500}{25}$
 = 0.0567 mol
 mass Fe = 0.0567×55.8
 = 3.164 g
 purity Fe = $\frac{3.164}{5.68} \times 100$
 = 55.7%



mol MnO_4^- = $0.0120 \times \frac{21.1}{1000}$
 = $2.532 \times 10^{-4} \text{ mol}$
 mol V^{2+} in 50 cm^3 = $\frac{5}{3} \times 2.532 \times 10^{-4}$
 = $4.22 \times 10^{-4} \text{ mol}$
 mol V^{2+} in 500 cm^3 = $4.22 \times 10^{-4} \times \frac{500}{50}$
 = $4.22 \times 10^{-3} \text{ mol}$
 mass V = $50.9 \times 4.22 \times 10^{-3}$
 = 0.2148 g
 purity = $\frac{0.2148}{0.382} \times 100$
 = 56.2%

3. Moles of Fe^{2+} in solution A

mol MnO_4^- = $0.0200 \times \frac{23.55}{1000}$
 = $4.71 \times 10^{-4} \text{ mol}$
 mol Fe^{2+} in 25 cm^3 = $5 \times 4.71 \times 10^{-4}$
 = $2.355 \times 10^{-3} \text{ mol}$
 mol Fe^{2+} in 500 cm^3 = $2.355 \times 10^{-3} \times \frac{500}{25}$
 = 0.0471 mol

Moles of Fe^{2+} in solution made from solution A

mol MnO_4^- = $0.0200 \times \frac{28.93}{1000}$
 = $5.786 \times 10^{-4} \text{ mol}$
 mol Fe^{2+} in 25 cm^3 = $5 \times 5.786 \times 10^{-4}$
 = $2.893 \times 10^{-3} \text{ mol}$
 mol Fe^{2+} in 200 cm^3 = $2.893 \times 10^{-3} \times \frac{200}{25}$
 = 0.02314 mol
 mol Fe^{2+} and Fe^{3+} in A = $0.02314 \times \frac{500}{100}$
 = 0.11572 mol

% Fe which is Fe^{2+} = $\frac{0.0471}{0.11572} \times 100$
 = 40.7%

COPYRIGHT
PROTECTED







$$\begin{aligned}
 4. \quad \text{mol S}_2\text{O}_3^{2-} &= 0.01 \times \frac{31.55}{1000} \\
 &= 3.155 \times 10^{-4} \text{ mol} \\
 \text{mol I}_2 \text{ in } 20 \text{ cm}^3 &= \frac{3.155 \times 10^{-4}}{2} \\
 &= 1.5775 \times 10^{-4} \text{ mol} \\
 \text{mol I}_2 \text{ in } 500 \text{ cm}^3 &= \frac{500}{20} \times 1.5775 \times 10^{-4} \\
 &= 0.003944 \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{mol BaFeO}_4 &= 0.676 \div (137.3 + 55.8 \times 4) \\
 &= 2.629 \times 10^{-3}
 \end{aligned}$$

	I ₂	FeO ₄ ²⁻
mol ratio	3.944 × 10 ⁻³	2.629 × 10 ⁻³
whole number	$\frac{3.944 \times 10^{-3}}{2.629 \times 10^{-3}}$	$\frac{2.629 \times 10^{-3}}{2.629 \times 10^{-3}}$
mol ratio	1.5	1
	3	2

$$\text{ratio FeO}_4^{2-} : \text{I}_2 = 3 : 2$$



$$\begin{aligned}
 5. \quad \text{mol NH}_4\text{FeSO}_4 &= 0.109 \times \frac{24.50}{1000} \\
 &= 2.6705 \times 10^{-3} \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{mol Cr}_2\text{O}_7^{2-} \text{ remaining} &= \frac{2.6705 \times 10^{-3}}{2} \\
 &= 1.33525 \times 10^{-3} \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{original mol Cr}_2\text{O}_7^{2-} &= 0.115 \times \frac{25}{1000} \\
 &= 2.875 \times 10^{-3} \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{mol Cr}_2\text{O}_7^{2-} \text{ used up} &= 2.875 \times 10^{-3} - 1.33525 \times 10^{-3} \\
 &= 1.53975 \times 10^{-3} \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{mol CH}_3\text{CH}_2\text{OH in sample} &= \frac{3}{2} \times 1.53975 \times 10^{-3} \\
 &= 2.309625 \times 10^{-3} \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{mol CH}_3\text{CH}_2\text{OH in } 500 \text{ cm}^3 &= 2.309625 \times 10^{-3} \times \frac{500}{20} \\
 &= 0.057740625 \text{ mol}
 \end{aligned}$$

$$\begin{aligned}
 \text{conc CH}_3\text{CH}_2\text{OH in wine} &= 0.057740625 \div \frac{100}{1000} \\
 &= 0.57740625 \text{ mol dm}^{-3}
 \end{aligned}$$

**COPYRIGHT
PROTECTED**



6. Overall equation: $\text{Mo}^{5+} + \text{Ce}^{4+} \rightarrow \text{Ce}^{3+} + \text{Mo}^{6+}$

$$\begin{aligned} \text{mol Ce}^{3+} &= 0.103 \times \frac{20.05}{1000} \\ &= 2.0652 \times 10^{-3} \text{ mol} \end{aligned}$$

$$\text{mol Mo}^{5+} \text{ in } 50.0 \text{ cm}^3 = 2.0652 \times 10^{-3} \text{ mol}$$

$$\begin{aligned} \text{mol Mo}^{5+} \text{ in } 200 \text{ cm}^3 &= 2.0652 \times 10^{-3} \times \frac{200}{50} \\ &= 8.26 \times 10^{-3} \text{ mol} \end{aligned}$$

$$\begin{aligned} \text{r.f.m. A} &= 2.00 \div 8.26 \times 10^{-3} \\ &= 242.1 \end{aligned}$$


$$\begin{aligned} \text{mol A in } 1.5 \text{ g} &= \frac{1.5}{242.1} \\ &= 6.196 \times 10^{-3} \text{ mol} \end{aligned}$$

$$\begin{aligned} \text{mass H}_2\text{O removed} &= 1.5 - 1.28 \\ &= 0.22 \text{ g} \end{aligned}$$

$$\begin{aligned} \text{mol H}_2\text{O removed} &= 0.22 \div 18 \\ &= 0.0122 \text{ mol} \end{aligned}$$

	$\text{Na}_2\text{MoO}_x \cdot n\text{H}_2\text{O}$	H_2O
mol ratio	6.196×10^{-3}	0.0122
whole number	6.196×10^{-3}	0.0122
mol ratio	$\frac{6.196 \times 10^{-3}}{6.196 \times 10^{-3}}$	$\frac{0.0122}{6.196 \times 10^{-3}}$
	1	2

$$n = 2$$



$$\text{r.f.m. due to O}_x = 242.1 - (2 \times 23 + 96) = 2 \times (2 + 16)$$

$$\begin{aligned} x &= 6 \\ &= 4 \end{aligned}$$




INSPECTION COPY

COPYRIGHT
PROTECTED



Exam-Style Questions: Answers

Question	Marking Guidance																				
1a	starts with an upwards curve ends at 40 cm³ vertical section at 25 cm³ 																				
1bi	<table border="1"> <thead> <tr> <th></th><th>Rough</th><th>1</th><th>2</th><th>3</th></tr> </thead> <tbody> <tr> <td>Start volume (cm³)</td><td>0.00</td><td>0.10</td><td>0.05</td><td>0.00</td></tr> <tr> <td>End volume (cm³)</td><td>33.35</td><td>31.15</td><td>31.15</td><td>31.10</td></tr> <tr> <td>Volume of NaOH (cm³)</td><td>33.35</td><td>31.05</td><td>31.10</td><td>31.10</td></tr> </tbody> </table> Mean titre = $\frac{31.05 + 31.10 + 31.10}{3}$ = 31.08 cm³		Rough	1	2	3	Start volume (cm ³)	0.00	0.10	0.05	0.00	End volume (cm ³)	33.35	31.15	31.15	31.10	Volume of NaOH (cm ³)	33.35	31.05	31.10	31.10
	Rough	1	2	3																	
Start volume (cm ³)	0.00	0.10	0.05	0.00																	
End volume (cm ³)	33.35	31.15	31.15	31.10																	
Volume of NaOH (cm ³)	33.35	31.05	31.10	31.10																	
1bii	mol NaOH = $0.100 \times \frac{31.08}{1000}$ = 3.108×10^{-3} mol mol organic acid = 3.108×10^{-3} mol conc organic acid = $\frac{3.108 \times 10^{-3} \times 25}{1000}$ = 0.024 mol dm⁻³																				
1c	$K_a = \frac{[H^+][A^-]}{[HA]}$ $[H^+] = [A^-]$ $K_a = \frac{[H^+]^2}{[HA]}$ $[H^+] = \sqrt{K_a \times [HA]}$ = $\sqrt{7.41 \times 10^{-4} \times 0.150}$ = 0.01054 mol dm⁻³ $K_w = [H^+] \times [OH^-]$ $[OH^-] = \frac{K_w}{[H^+]}$ = $\frac{1 \times 10^{-14}}{0.01054}$ = 9.48×10^{-13} mol dm⁻³																				

INSPECTION COPY

COPYRIGHT
PROTECTED



Question	Marking guidance
1di	$\text{H}_3\text{C}_6\text{H}_5\text{O}_7 \rightleftharpoons \text{H}_2\text{C}_6\text{H}_5\text{O}_7^- + \text{H}^+$ $\text{H}_2\text{C}_6\text{H}_5\text{O}_7^-$ reacts with HCl / H^+ Equilibrium position shifts left to reduce the concentration of H^+ again / H^+ concentration
1dii	$[\text{H}^+] = 10^{-5}$ $= 1 \times 10^{-5} \text{ mol dm}^{-3}$ $K_a = 7.41 \times 10^{-4}$ $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$ assume $[\text{NaOH}] = 0.1$ $[\text{H}^+]_{\text{initial}} - [\text{NaOH}] = \frac{[\text{H}^+][\text{NaOH}]}{K_a}$ $[\text{HA}]_{\text{initial}} = \frac{[\text{H}^+][\text{NaOH}]}{K_a} + [\text{NaOH}]$ $= \frac{1 \times 10^{-5} \times 0.1}{7.41 \times 10^{-4}} + 0.1$ $= 0.101 \text{ mol dm}^{-3}$

Question	Marking guidance
2	Stage 1: Acid is in excess 'Hydrochloric acid must be in excess.' $\text{mol HCl} = 0.150 \times \frac{100}{1000}$ $= 0.015 \text{ mol}$ 2 : 1 ratio $\text{HCl} : \text{CaCO}_3$ max. mol $\text{CaCO}_3 = \frac{0.015}{2}$ $= 7.5 \times 10^{-3} \text{ mol}$ $\text{mass CaCO}_3 = 7.5 \times 10^{-3} \times (40.1 + 12 + 3 \times 16)$ $= 0.751 \text{ g}$ Stage 2: Experimental method Measure 100 cm^3 HCl (with a pipette) and add the mass of calcite weighed into a conical flask. Titrate: standard solution / $0.100 \text{ mol dm}^{-3}$ / $0.200 \text{ mol dm}^{-3}$ NaOH in a standard solution and record the volume when the indicator changes. Stage 3: Calculate mol CaCO_3 $\text{mol NaOH} = \frac{\text{volume of NaOH in cm}^3}{1000} \times [\text{NaOH}]$ AND $\text{mol HCl} = 7.5 \times 10^{-3} - \text{mol NaOH}$ AND $\text{mol CaCO}_3 = \frac{\text{mol HCl}}{2}$ Stage 4: Calculate purity $\text{mass CaCO}_3 = \text{mol CaCO}_3 \times 100.1$ AND $\text{Purity} = \frac{\text{mass CaCO}_3}{\text{mass of solid}} \times 100$

**COPYRIGHT
PROTECTED**



Question	Marking Guidance
3a	$\text{Cu}^{2+} + \text{e}^{-} \rightarrow \text{Cu}^{+}$
3b	Because it might react with sodium thiosulfate, causing the result / titre / percentage by mass to be inaccurate
3c	Indicator
3d	$\text{mol Na}_2\text{S}_2\text{O}_3 = \frac{37.20}{1000} \times 0.01$ $= 0.000372 \text{ mol}$ $\text{mol I}_2 = \frac{0.000372}{2}$ $= 0.000186 \text{ mol}$ $\text{mol I}_2 \text{ in } 250 \text{ cm}^3 = 0.000186 \times \frac{250}{25}$ $= 0.00186 \text{ mol}$ $\text{mol Cu in ring} = 0.00186 \times 2$ $= 0.00372 \text{ mol}$ $\text{Mass Cu in ring} = 0.00372 \times 63.5$ $= 0.2362 \text{ g}$ $\% \text{ by mass Cu} = \frac{0.2362}{0.945} \times 100$ $= 25.0 \%$

Question	Marking Guidance
4	Full equation: $2\text{MnO}_4^{-} + 5\text{SO}_3^{2-} + 6\text{H}^{+} \rightarrow 2\text{Mn}^{2+} + 5\text{SO}_4^{2-} + 3\text{H}_2\text{O}$ $\text{mol MnO}_4^{-} = \frac{0.0200}{1000} \times \frac{1}{2}$ $= 1.0 \times 10^{-4} \text{ mol}$ $\text{mol SO}_3^{2-} = \frac{1.0}{2} \times 4.92 \times 10^{-4}$ $= 1.23 \times 10^{-3} \text{ mol}$ $\text{mol SO}_3^{2-} \text{ in } 200 \text{ cm}^3 = 1.23 \times 10^{-3} \times \frac{200}{25}$ $= 9.84 \times 10^{-3} \text{ mol}$ $\text{r.f.m. M}_2\text{SO}_3 = \frac{1.24}{9.84 \times 10^{-3}}$ $= 126.0$ $\text{r.f.m. M} = \frac{126.0 - (32.1 + 3 \times 16)}{2}$ $= 23$ $\text{M}^{+} = \text{Na}^{+}$

COPYRIGHT
PROTECTED



Quick Quiz

1. H^+
2. Neutralisation
3. hydrochloric acid + potassium hydroxide $\rightarrow$ potassium chloride
 $\text{HCl} \quad \text{KOH} \quad \rightarrow \quad \text{KCl} \quad +$
4. Neutral
5. pH meter
(universal indicator)
6. $\frac{3}{9} = 0.33 \text{ dm}^{-3}$
7. Any three from:
 - burette
 - pipette
 - beaker
 - conical flask
 - clamp / clamp stand / retort stand
 - funnel
 - white tile
 - measuring cylinder
8. Strong acids / sulfuric acid / **solution 1** dissociate(s) more easily than weak acids /
9. a) $\text{H}_2\text{SO}_4 + 2\text{NaOH} \rightarrow \text{Na}_2\text{SO}_4 + 2\text{H}_2\text{O}$
b) 0.24 mol
10. a) Fe
b) C
c) O

INSPECTION COPY

COPYRIGHT
PROTECTED

